


FORM PTO-1390 (REV. 11-2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER A0000102-01-SMH	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				U.S. APPLICATION NO. (If known, see 37 CFR 1.5) <b>10/031037</b>	
INTERNATIONAL APPLICATION NO. PCT/US00/18345		INTERNATIONAL FILING DATE 5 July 2000		PRIORITY DATE CLAIMED July 16, 1999	
TITLE OF INVENTION Method for Treating Chronic Pain Using Mek Inhibitors					
APPLICANT(S) FOR DO/EO/US BARRETT, Stephen Douglas, BRIDGES, Alexander James, DIXON, Alistair, LEE, Kevin, PINNOCK, Robert Denham, TECLE, Haile					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
<p>1. <input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371.</p> <p>2. <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371.</p> <p>3. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.</p> <p>4. <input type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31).</p> <p>5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2))</p> <p style="margin-left: 20px;">a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau).</p> <p style="margin-left: 20px;">b. <input checked="" type="checkbox"/> has been communicated by the International Bureau.</p> <p style="margin-left: 20px;">c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</p> <p>6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).</p> <p style="margin-left: 20px;">a. <input type="checkbox"/> is attached hereto.</p> <p style="margin-left: 20px;">b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4).</p> <p>7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))</p> <p style="margin-left: 20px;">a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau).</p> <p style="margin-left: 20px;">b. <input type="checkbox"/> have been communicated by the International Bureau.</p> <p style="margin-left: 20px;">c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</p> <p style="margin-left: 20px;">d. <input type="checkbox"/> have not been made and will not be made.</p> <p>8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).</p> <p>9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).</p> <p>10. <input type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).</p> <p><b>Items 11 to 20 below concern document(s) or information included:</b></p> <p>11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.</p> <p>12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.</p> <p>13. <input type="checkbox"/> A <b>FIRST</b> preliminary amendment.</p> <p>14. <input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment.</p> <p>15. <input type="checkbox"/> A substitute specification.</p> <p>16. <input type="checkbox"/> A change of power of attorney and/or address letter.</p> <p>17. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.</p> <p>18. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4).</p> <p>19. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).</p> <p>20. <input checked="" type="checkbox"/> Other items or information:  International Search Report (PCT/ISA/210)  International Preliminary Examination Report (PCT/IPEA/409)  Certificate of Mailing by Express Mail</p>					

Express Mail No. EF378126665US  
PD-A0000102-01-SMH

U.S. APPLICATION NO. (if known, see 37 CFR 1.53) <b>10/031037</b>		INTERNATIONAL APPLICATION NO. PCT/US00/18345		ATTORNEY'S DOCKET NUMBER A0000102-01-SMH	
<b>21. <input checked="" type="checkbox"/> The following fees are submitted:</b> <b>BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):</b> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO ..... <b>\$1000.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... <b>\$860.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... <b>\$710.00</b>  International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... <b>\$690.00</b>  International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) ..... <b>\$100.00</b>  <b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b>				<b>CALCULATIONS PTO USE ONLY</b>	
Surcharge of <b>\$130.00</b> for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$ 860.00	
Surcharge of <b>\$130.00</b> for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$ 130.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$	
Total claims	32 - 20 =	12	x <b>\$18.00</b>	\$ 216.00	
Independent claims	1 - 3 =		x <b>\$80.00</b>	\$ 0.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ <b>\$270.00</b>	\$	
<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$ 1,206.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	
<b>SUBTOTAL =</b>				\$	
Processing fee of <b>\$130.00</b> for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
<b>TOTAL NATIONAL FEE =</b>				\$ 1,206.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). <b>\$40.00</b> per property +				\$	
<b>TOTAL FEES ENCLOSED =</b>				\$ 1,206.00	
				Amount to be refunded:	\$
				charged:	\$
a. <input type="checkbox"/> A check in the amount of \$ _____ to cover the above fees is enclosed.					
b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. <u>23-0455</u> in the amount of \$ <u>1,206.00</u> to cover the above fees. A duplicate copy of this sheet is enclosed.					
c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>23-0455</u> . A duplicate copy of this sheet is enclosed.					
d. <input type="checkbox"/> Fees are to be charged to a credit card. <b>WARNING:</b> Information on this form may become public. <b>Credit card information should not be included on this form.</b> Provide credit card information and authorization on PTO-2038.					
<b>NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.</b>					
SEND ALL CORRESPONDENCE TO: Suzanne M. Harvey Warner-Lambert Company Patent Department 2800 Plymouth Road Ann Arbor, Michigan 48105 Telephone: (734) 622-2658 Facsimile: (734) 622-1553					
				 SIGNATURE	
				Suzanne M. Harvey NAME	
				<u>42,640</u> REGISTRATION NUMBER	

**METHOD FOR TREATING CHRONIC PAIN**  
**USING MEK INHIBITORS**

5 BACKGROUND

The invention features a method for treating chronic pain using MEK inhibitors. Chronic pain includes neuropathic pain, and chronic inflammatory pain.

10 Abnormality anywhere in a nerve pathway disrupts nerve signals, which in turn are abnormally interpreted in the brain, causing neuropathic pain. Neuropathic pain may be, for example, a deep ache, a burning sensation, or hypersensitivity to touch. Diseases or conditions associated with neuropathic pain include, without limitation, diabetic neuropathy, causalgia, plexus  
15 avulsion, neuroma, vasculitis, crush injury, viral infections (e.g., herpes virus infection or HIV), constriction injury, tissue injury, nerve injury from the periphery to the central nervous system, limb amputation, hypothyroidism, uremia, chronic alcoholism, post-operative pain, arthritis, back pain, and vitamin deficiencies.

20 Infections such as herpes zoster (shingles) can cause nerve inflammation and produce postherpetic neuralgia, a chronic burning localized to the area of viral infection. Hyperalgesia is when an already noxious stimulus becomes more painful, and allodynia, when a previously non-noxious stimulus becomes painful (such as contact of clothing or a breeze). Reflex  
25 sympathetic dystrophy is accompanied by swelling and sweating or changes in local blood flow, tissue atrophy, or osteoporosis. Causalgia, including severe burning pain and swelling, sweating, and changes in blood flow, may follow an injury or disease of a major nerve such as the sciatic nerve. Some types of chronic low back pain can have a neuropathic component (e.g.,  
30 sciatica, postpoliomyelitis and CPRM). Neuropathic pain may also be induced by cancer or chemotherapy.

Neuropathic pain is currently treated with anticonvulsants such as carbamazepine and antidepressants such as amitriptyline. NSAIDs and opioids generally have little effect (*Fields et al 1994 Textbook of Pain p 991-996 (pub: Churchill Livingstone), James & Page 1994*

5 *J.Am.Pediatr.Med.Assoc, 8: 439-447, Galer, 1995 Neurology 45 S17-S25.*


Neuropathic conditions that have been treated with gabapentin include: postherpetic neuralgia, postpoliomyelitis, CPRM, HIV-related neuropathy, trigeminal neuralgia, and reflex sympathetic dystrophy (RSD).

The generally weak efficacy of antiinflammatory agents suggests that the  
10 mechanism for chronic pain is separate from hyperalgesia.

### SUMMARY OF THE INVENTION

The invention features a method for treating chronic pain, which  
15 method includes the step of administering a composition including a MEK inhibitor to a patient in need of such treatment. Chronic pain includes neuropathic pain, idiopathic pain, and pain associated with vitamin deficiencies, uremia, hypothyroidism post-operative pain, arthritis, back pain, and chronic alcoholism. The invention also features compositions as  
20 disclosed, formulated for the treatment of chronic pain. Such a composition may include one or more MEK inhibitor compounds having a structure disclosed in patent applications USSN 60/115,873, filed January 13, 1999, PCT/US99/30483, international filing date December 21, 1999.


Examples of MEK inhibitors include a compound having the formula (I)  
25 below:




(I)




- In formula (I), W is  $OR_1$ ,  $NR_2OR_1$ ,  $NR_AR_B$ ,  $NR_2NR_AR_B$ ,  $O(CH_2)_{2-4}NR_AR_B$ , or  $NR_2(CH_2)_{2-4}NR_AR_B$ .  $R_1$  is H, C<sub>1-8</sub> alkyl, C<sub>3-8</sub> alkenyl, C<sub>3-8</sub> alkynyl, C<sub>3-8</sub> cycloalkyl, phenyl, (phenyl)C<sub>1-4</sub> alkyl, (phenyl)C<sub>3-4</sub> alkenyl, (phenyl)C<sub>3-4</sub> alkynyl, (C<sub>3-8</sub> cycloalkyl)C<sub>1-4</sub> alkyl, (C<sub>3-8</sub> cycloalkyl)C<sub>3-4</sub> alkenyl, (C<sub>3-8</sub> cycloalkyl)C<sub>3-4</sub> alkynyl, C<sub>3-8</sub> heterocyclic radical, (C<sub>3-8</sub> heterocyclic radical)C<sub>1-4</sub> alkyl, (C<sub>3-8</sub> heterocyclic radical)C<sub>3-4</sub> alkenyl, (C<sub>3-8</sub> heterocyclic radical)C<sub>3-4</sub> alkynyl or  $(CH_2)_{2-4}NR_CR_D$ .  $R_2$  is H, C<sub>1-4</sub> alkyl, phenyl, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> heterocyclic radical, or (C<sub>3-6</sub> cycloalkyl) methyl.  $R_A$  is H, C<sub>1-6</sub> alkyl, C<sub>3-8</sub> alkenyl, C<sub>3-8</sub> alkynyl, C<sub>3-8</sub> cycloalkyl, phenyl, (C<sub>3-8</sub> cycloalkyl)C<sub>1-4</sub> alkyl, (C<sub>3-8</sub> cycloalkyl)C<sub>3-4</sub> alkenyl, (C<sub>3-8</sub> cycloalkyl)C<sub>3-4</sub> alkynyl, C<sub>3-8</sub> heterocyclic radical, (C<sub>3-8</sub> heterocyclic radical)C<sub>1-4</sub> alkyl, (aminosulfonyl)phenyl, [(aminosulfonyl)phenyl]C<sub>1-4</sub> alkyl, (aminosulfonyl)C<sub>1-6</sub> alkyl, (aminosulfonyl)C<sub>3-6</sub> cycloalkyl, [(aminosulfonyl)C<sub>3-6</sub> cycloalkyl]C<sub>1-4</sub> alkyl, or  $(CH_2)_{2-4}NR_CR_D$ .  $R_B$  is H, C<sub>1-8</sub> alkyl, C<sub>3-8</sub> alkenyl, C<sub>3-8</sub> alkynyl, C<sub>3-8</sub> cycloalkyl, or phenyl.
- Q is one of the following formulae (i) – (iii):



(i)




(ii)




(iii)


$R_3$  is H or F;  $R_4$  is halo,  $\text{NO}_2$ ,  $\text{SO}_2\text{NR}_\text{O}(\text{CH}_2)_{2-4}\text{NR}_\text{E}R_\text{F}$ ,  $\text{SO}_2\text{NR}_\text{E}R_\text{F}$  or  $(\text{CO})\text{T}$ . T is  $\text{C}_{1-8}$  alkyl,  $\text{C}_{3-8}$  cycloalkyl,  $(\text{NR}_\text{E}R_\text{F})\text{C}_{1-4}$  alkyl,  $\text{OR}_\text{F}$ ,  $-\text{NR}_\text{O}(\text{CH}_2)_{2-4}\text{NR}_\text{E}R_\text{F}$ , or  $\text{NR}_\text{E}R_\text{F}$ ; Z is one of the following formulae (iv) – (viii):




(iv)




(v)



(vi)



(vii)



(viii)

- 5
- 10 One of  $R_5$  and  $R_6$  is H or methyl and the other of  $R_5$  and  $R_6$  is H,  $\text{C}_{1-6}$  alkyl,  $\text{C}_{2-6}$  alkenyl,  $\text{C}_{2-6}$  alkynyl, phenyl, benzyl, or  $-\text{M}-\text{E}-\text{G}$ . M is O, CO,  $\text{SO}_2$ ,  $\text{NR}_\text{J}$ ,  $(\text{CO})\text{NR}_\text{H}$ ,  $\text{NR}_\text{H}(\text{CO})$ ,  $\text{NR}_\text{H}(\text{SO}_2)$ ,  $(\text{SO}_2)\text{NR}_\text{H}$ , or  $\text{CH}_2$ . E is  $(\text{CH}_2)_{1-4}$  or  $(\text{CH}_2)_\text{m}$   $\text{O}(\text{CH}_2)_\text{p}$  where  $1 \leq (\text{each of m and p}) \leq 3$  and  $2 \leq (m + p) \leq 4$ ; or E is absent. G is  $\text{R}_\text{K}$ ,  $\text{OR}_\text{I}$  or  $\text{NR}_\text{J}\text{R}_\text{K}$ , provided that if  $p = 1$ , then G is H.  $R_7$  is H,  $\text{C}_{1-4}$  alkyl,
- 15  $\text{C}_{2-4}$  alkenyl,  $\text{C}_{2-4}$  alkynyl,  $\text{C}_{3-6}$  cycloalkyl, phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl,  $(\text{CH}_2)_{1-2}\text{Ar}$ , where Ar is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl,  $\text{SO}_2\text{NR}_\text{H}(\text{CH}_2)_{2-4}\text{NR}_\text{J}\text{R}_\text{K}$ ,  $(\text{CO})(\text{CH}_2)_{2-4}\text{NR}_\text{J}\text{R}_\text{K}$  or  $(\text{CO})\text{NR}_\text{H}(\text{CH}_2)_{2-4}\text{NR}_\text{J}\text{R}_\text{K}$ .  $X_1$  is O, S,  $\text{NR}_\text{8}$ , or  $\text{CHR}_\text{9}$ ;  $X_2$  is O, S, or  $\text{CHR}_\text{9}$ ; and  $X_3$  is O or S. In one embodiment, if  $X_1$  or  $X_2$  is  $\text{CHR}_\text{9}$ , the disclosed compound may also be a
- 20 tautomerized indole.  $R_8$  is H,  $\text{C}_{1-4}$  alkyl, phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl,  $(\text{CH}_2)_{1-2}\text{Ar}$ , where Ar is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl,  $\text{C}_{2-4}$  alkenyl,

C<sub>2-4</sub> alkynyl, C<sub>3-6</sub> cycloalkyl, or (C<sub>2-4</sub> alkyl)NR<sub>L</sub>R<sub>M</sub> provided R<sub>7</sub> and R<sub>8</sub> together have no more than 14 carbon atoms, exclusive of R<sub>L</sub>, R<sub>M</sub>, R<sub>J</sub> and R<sub>K</sub>.

R<sub>G</sub> is C<sub>1-4</sub> alkyl, phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, C<sub>3-4</sub> alkenyl, C<sub>3-4</sub> alkynyl, C<sub>3-6</sub> cycloalkyl, (CO)OR<sub>P</sub>, (C<sub>2-4</sub> alkyl)NR<sub>L</sub>R<sub>M</sub>, (CO)NR<sub>N</sub>(CH<sub>2</sub>)<sub>2</sub>-

5 4NR<sub>L</sub>R<sub>M</sub>, (CO)NR<sub>L</sub>R<sub>M</sub>, (CO)(CH<sub>2</sub>)<sub>2-4</sub>-NR<sub>L</sub>R<sub>M</sub>, or (CH<sub>2</sub>)<sub>1-2</sub>Ar, where Ar is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl. R<sub>9</sub> is C<sub>1-4</sub> alkyl, phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, C<sub>3-6</sub> cycloalkyl, (CO)OR<sub>P</sub>, (C<sub>2-4</sub> alkyl)NR<sub>L</sub>R<sub>M</sub>, (CO)NR<sub>N</sub>(CH<sub>2</sub>)<sub>2-4</sub>NR<sub>L</sub>R<sub>M</sub>, (CO)NR<sub>L</sub>R<sub>M</sub>, (CO)(CH<sub>2</sub>)<sub>2-4</sub>-NR<sub>L</sub>R<sub>M</sub>, or (CH<sub>2</sub>)<sub>1-2</sub>Ar', where Ar' is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl.

10 R<sub>P</sub> is H, C<sub>1-6</sub> alkyl, phenyl, C<sub>3-4</sub> alkenyl, C<sub>3-4</sub> alkynyl, C<sub>3-6</sub> cycloalkyl, or (CH<sub>2</sub>)<sub>2-4</sub>NR<sub>L</sub>R<sub>M</sub>; R<sub>10</sub> is H, methyl, halo, or NO<sub>2</sub>; R<sub>11</sub> is H, methyl, halo, or NO<sub>2</sub>. Each of R<sub>C</sub>, R<sub>D</sub>, R<sub>E</sub>, R<sub>F</sub>, R<sub>I</sub>, R<sub>J</sub>, R<sub>K</sub>, R<sub>L</sub> and R<sub>M</sub> is independently selected from H, C<sub>1-4</sub> alkyl, C<sub>3-4</sub> alkenyl, C<sub>3-4</sub> alkynyl, C<sub>3-6</sub> cycloalkyl, and phenyl; each of NR<sub>C</sub>R<sub>D</sub>, NR<sub>E</sub>R<sub>F</sub>, NR<sub>J</sub>R<sub>K</sub>, and NR<sub>L</sub>R<sub>M</sub> can also independently be morpholinyl, 15 piperazinyl, pyrrolidinyl, or piperadiny. Each of R<sub>H</sub>, R<sub>N</sub>, and R<sub>O</sub> is independently H, methyl, or ethyl. Finally, each hydrocarbon radical or heterocyclic radical above is optionally substituted with between 1 and 3 substituents independently selected from halo, C<sub>1-4</sub> alkyl, C<sub>3-6</sub> cycloalkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl, phenyl, hydroxyl, amino, (amino)sulfonyl, and NO<sub>2</sub>, 20 wherein each substituent alkyl, cycloalkyl, alkenyl, alkynyl or phenyl is in turn optionally substituted with between 1 and 3 substituents independently selected from halo, C<sub>1-2</sub> alkyl, hydroxyl, amino, and NO<sub>2</sub>. In addition to the above compounds, the invention also provides a pharmaceutically-acceptable salt or C<sub>1-7</sub> ester thereof.

25 Preferred embodiments of the invention include methods using one or more of the following compounds:

(a) said MEK inhibitor has a structure selected from: 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1*H*-benzoimidazole-5-carboxylic acid

cyclopropylmethoxy-amide; 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-6,7-

30 dihydro-1*H*-benzoimidazole-5-carboxylic acid (hydrochloride); 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1*H*-benzoimidazole-5-carboxylic acid; 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-3*H*-benzoimidazole-5-carboxylic acid

(2-hydroxy-ethoxy)-amide; 6-(2-chloro-4-iodo-phenylamino)-7-fluoro-1*H*-benzoimidazole-5-carboxylic acid; and 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1*H*-benzoimidazole-5-carboxylic acid pentafluorophenyl ester; and (b) said MEK inhibitor has a structure selected from: 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1*H*-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide; and 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-3*H*-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide.

The invention also relates to a pharmaceutical composition including (a) a benzoheterocycle (e.g., of formula I) and (b) a pharmaceutically-acceptable carrier.

### BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 is a bar graph representing the paw withdrawal threshold (PWT) in grams as a function of time in days. The empty, cross-hatched, and single-hatched bars are vehicle, PD 198306, and pregabalin, respectively. The arrows indicate time of drug administration (30 mg/kg, p.o.).

FIG. 2. is a bar graph representing the force required in grams to elicit paw withdrawal using von Frey hair filaments as a function of time in days. Baseline (BL) measurements were taken before treatment. Animals received a single p.o. administration of PD 198306 (3-30mg/kg), or pregabalin (30mg/kg) and withdrawal thresholds were re-assessed 1h after treatment. Treatments were repeated twice a day for two days. Results are expressed median  $\pm$  1<sup>st</sup> and 3<sup>rd</sup> quartiles. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  significantly different from vehicle treated animals (Mann-Whitney t test;  $n = 7-8$ ).

FIG. 3. is a bar graph representing the force required in grams to elicit paw withdrawal using von Frey hair filaments as a function of time in days. Baseline (BL) measurements were taken before treatment. Animals received a single p.o. administration of PD 198306 (3-30mg/kg), or pregabalin (30mg/kg) and withdrawal thresholds were re-assessed 1h after treatment.

Treatments were repeated twice a day for two days. Results are expressed median  $\pm$  1<sup>st</sup> and 3<sup>rd</sup> quartiles. \*\*P<0.01 significantly different from vehicle treated animals (Mann-Whitney t test; n=6).

5           FIG. 4. is a bar graph representing the force required in grams to elicit paw withdrawal using von Frey hair filaments as a function of time in days. Baseline (BL) measurements were taken before treatment. Animals received a single i.t. administration of PD 198306 (1-30 $\mu$ g/10 $\mu$ l), or pregabalin (100 $\mu$ g/10 $\mu$ l) and withdrawal thresholds were re-assessed at 30min, 1h and  
10   2h after treatment. Results are expressed median  $\pm$  1<sup>st</sup> and 3<sup>rd</sup> quartiles. \*P<0.05, \*\*\*P<0.001 significantly different from vehicle treated animals (Mann-Whitney t test; n=7-9).

15           FIG. 5. is a bar graph representing the force required in grams to elicit paw withdrawal using von Frey hair filaments as a function of time in days. Baseline (BL) measurements were taken before treatment. Animals received a single i.t. administration of PD 198306 (1-30 $\mu$ g/10 $\mu$ l), or pregabalin (100 $\mu$ g/10 $\mu$ l) and withdrawal thresholds were re-assessed at 30min, 1h and  
20   2h after treatment. Results are expressed median  $\pm$  1<sup>st</sup> and 3<sup>rd</sup> quartiles. \*P<0.05, \*\*P<0.01, \*\*\*P<0.001 significantly different from vehicle treated animals (Mann-Whitney t test; n=6-8).

25           FIG. 6 is a bar graph representing the force required in grams to elicit paw withdrawal using von Frey hair filaments as a function of time in days . Animals received a single intraplantar (i.pl.) administration of PD 198306 (3mg/100 $\mu$ l), or an intrathecal injection of PD 198306 (30 $\mu$ g/10 $\mu$ l) and withdrawal thresholds were re-assessed 1h after treatment. Results are expressed median  $\pm$  1<sup>st</sup> and 3<sup>rd</sup> quartiles. \*\*P<0.01 significantly different from vehicle treated animals (Mann-Whitney t test; n=6-9).

30           FIG. 7. is a bar graph representing the force required in grams to elicit paw withdrawal using von Frey hair filaments as a function of time in days.

Animals received a single intraplantar (i.pl.) administration of PD 198306 (3mg/100 $\mu$ l), or an intrathecal injection of PD 198306 (30 $\mu$ g/10 $\mu$ l) and withdrawal thresholds were re-assessed 1h after treatment. Results are expressed median  $\pm$  1<sup>st</sup> and 3<sup>rd</sup> quartiles. \*\*P<0.01 significantly different from vehicle treated animals (Mann-Whitney t test; n=6).

FIG. 8 is a bar graph representing the force required in grams to elicit paw withdrawal using von Frey hair filaments. Baseline (BL) measurements were taken before treatment. Animals received a single i.t. administration of PD219622, PD297447, PD 184352, or PD 254552 (30 $\mu$ g/10 $\mu$ l), or pregabalin (100 $\mu$ g/10 $\mu$ l) and withdrawal thresholds were re-assessed at 30min, 1h and 2h after treatment. Results are expressed median  $\pm$  1<sup>st</sup> and 3<sup>rd</sup> quartiles. \*P<0.05, \*\*P<0.01, \*\*\*P<0.001 significantly different from vehicle treated animals (Mann-Whitney t test; n=7-8).

#### DETAILED DESCRIPTION

The compounds disclosed herein are pharmaceutically active, for example, they inhibit MEK. MEK enzymes are dual specificity kinases involved in, for example, immunomodulation, inflammation, and proliferative diseases such as cancer and restenosis.

Proliferative diseases are caused by a defect in the intracellular signaling system, or the signal transduction mechanism of certain proteins. Defects include a change either in the intrinsic activity or in the cellular concentration of one or more signaling proteins in the signaling cascade. The cell may produce a growth factor that binds to its own receptors, resulting in an autocrine loop, which continually stimulates proliferation. Mutations or overexpression of intracellular signaling proteins can lead to spurious mitogenic signals within the cell. Some of the most common mutations occur in genes encoding the protein known as Ras, a G-protein that is activated when bound to GTP, and inactivated when bound to GDP. The above-mentioned growth factor receptors, and many other mitogenic receptors, when

activated, lead to Ras being converted from the GDP-bound state to the GTP-bound state. This signal is an absolute prerequisite for proliferation in most cell types. Defects in this signaling system, especially in the deactivation of the Ras-GTP complex, are common in cancers, and lead to the signaling cascade below Ras being chronically activated.

Activated Ras leads in turn to the activation of a cascade of serine/threonine kinases. One of the groups of kinases known to require an active Ras-GTP for its own activation is the Raf family. These in turn activate MEK (e.g., MEK<sub>1</sub> and MEK<sub>2</sub>) which then activates MAP kinase, ERK (ERK<sub>1</sub> and ERK<sub>2</sub>). Activation of MAP kinase by mitogens appears to be essential for proliferation; constitutive activation of this kinase is sufficient to induce cellular transformation. Blockade of downstream Ras signaling, for example by use of a dominant negative Raf-1 protein, can completely inhibit mitogenesis, whether induced from cell surface receptors or from oncogenic Ras mutants. Although Ras is not itself a protein kinase, it participates in the activation of Raf and other kinases, most likely through a phosphorylation mechanism. Once activated, Raf and other kinases phosphorylate MEK on two closely adjacent serine residues, S<sup>218</sup> and S<sup>222</sup> in the case of MEK-1, which are the prerequisite for activation of MEK as a kinase. MEK in turn phosphorylates MAP kinase on both a tyrosine, Y<sup>185</sup>, and a threonine residue, T<sup>183</sup>, separated by a single amino acid.

This double phosphorylation activates MAP kinase at least 100-fold. Activated MAP kinase can then catalyze the phosphorylation of a large number of proteins, including several transcription factors and other kinases. Many of these MAP kinase phosphorylations are mitogenically activating for the target protein, such as a kinase, a transcription factor, or another cellular protein. In addition to Raf-1 and MEKK, other kinases activate MEK, and MEK itself appears to be a signal integrating kinase. Current understanding is that MEK is highly specific for the phosphorylation of MAP kinase. In fact, no substrate for MEK other than the MAP kinase, ERK, has been demonstrated to date and MEK does not phosphorylate peptides based on the MAP kinase phosphorylation sequence, or even phosphorylate denatured

MAP kinase. MEK also appears to associate strongly with MAP kinase prior to phosphorylating it, suggesting that phosphorylation of MAP kinase by MEK may require a prior strong interaction between the two proteins. Both this requirement and the unusual specificity of MEK are suggestive that it  
5 may have enough difference in its mechanism of action to other protein kinases that selective inhibitors of MEK, possibly operating through allosteric mechanisms rather than through the usual blockade of the ATP binding site, may be found.

The effect of the MEK inhibitor PD 198306 has been investigated in two  
10 animal models of neuropathic pain by assessing static allodynia with von Frey hairs.

Oral administration of PD 198306 (3-30mg/kg) had no effect in the model of chronic constriction injury of the sciatic nerve (CCI). However, after repeated administration (3 doses over two days) it had a transient effect in the diabetic neuropathy model (streptozocin). This may be due to disorders of the blood-  
15 brain barrier induced by the diabetic condition in these animals, thus allowing central action of the compound. Intrathecal administration of PD 198306 (1-30µg) dose-dependently blocked static allodynia in both the streptozocin and the CCI models of neuropathic pain, with minimum effective doses (MED) of 3 and  
20 10µg respectively. The highest dose used (30µg) totally blocked the maintenance of static allodynia, for up to 1h. Intraplantar administration of PD 198306 (3mg/100µl) at a dose 100-fold higher than the dose shown to be effective intrathecally (30µg/10µl) had no effect on static allodynia in either of the neuropathic pain models. This finding confirms the lack of effect seen after  
25 systemic administration and suggests a central site of action for the compound.

From this study we can suggest the use of MEK inhibitors as potential new therapeutic tools for chronic pain. The study of potential side-effects, especially related to memory, of future brain-penetrant MEK inhibitors will indicate the therapeutic window for this novel class of compounds in the  
30 treatment of pain.



## A. Terms

Certain terms are defined below and by their usage throughout this disclosure.

Alkyl groups include aliphatic (i.e., hydrocarbyl or hydrocarbon radical structures containing hydrogen and carbon atoms) with a free valence. Alkyl groups are understood to include straight chain and branched structures. Examples include methyl, ethyl, propyl, isopropyl, butyl, n-butyl, isobutyl, t-butyl, pentyl, isopentyl, 2,3-dimethylpropyl, hexyl, 2,3-dimethylhexyl, 1,1-dimethylpentyl, heptyl, and octyl. Cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

Alkyl groups can be substituted with 1, 2, 3 or more substituents which are independently selected from halo (fluoro, chloro, bromo, or iodo), hydroxy, amino, alkoxy, alkylamino, dialkylamino, cycloalkyl, aryl, aryloxy, arylalkyloxy, heterocyclic radical, and (heterocyclic radical)oxy. Specific examples include fluoromethyl, hydroxyethyl, 2,3-dihydroxyethyl, (2- or 3-furanyl)methyl, cyclopropylmethyl, benzyloxyethyl, (3-pyridinyl)methyl, (2- or 3-furanyl)methyl, (2-thienyl)ethyl, hydroxypropyl, aminocyclohexyl, 2-dimethylaminobutyl, methoxymethyl, N-pyridinylethyl, diethylaminoethyl, and cyclobutylmethyl.

Alkenyl groups are analogous to alkyl groups, but have at least one double bond (two adjacent  $sp^2$  carbon atoms). Depending on the placement of a double bond and substituents, if any, the geometry of the double bond may be *entgegen* (E), or *zusammen* (Z), *cis*, or *trans*. Similarly, alkynyl groups have at least one triple bond (two adjacent  $sp$  carbon atoms).

Unsaturated alkenyl or alkynyl groups may have one or more double or triple bonds, respectively, or a mixture thereof; like alkyl groups, unsaturated groups may be straight chain or branched, and they may be substituted as described both above for alkyl groups and throughout the disclosure by example.

Examples of alkenyls, alkynyls, and substituted forms include *cis*-2-butenyl, *trans*-2-butenyl, 3-butyne, 3-phenyl-2-propyne, 3-(2'-fluorophenyl)-2-propyne, 3-methyl(5-phenyl)-4-pentyne, 2-hydroxy-2-propyne, 2-methyl-2-propyne, 2-propenyl, 4-hydroxy-3-butyne, 3-(3-fluorophenyl)-2-propyne, and 2-methyl-2-

propenyl. In formula (I), alkenyls and alkynyls can be C<sub>2-4</sub> or C<sub>2-8</sub>, for example, and are preferably C<sub>3-4</sub> or C<sub>3-8</sub>.

More general forms of substituted hydrocarbon radicals include hydroxyalkyl, hydroxyalkenyl, hydroxyalkynyl, hydroxycycloalkyl, hydroxyaryl, and corresponding forms for the prefixes amino-, halo- (e.g., fluoro-, chloro-, or bromo-), nitro-, alkyl-, phenyl-, cycloalkyl- and so on, or combinations of substituents. According to formula (I), therefore, substituted alkyls include hydroxyalkyl, aminoalkyl, nitroalkyl, haloalkyl, alkylalkyl (branched alkyls, such as methylpentyl), (cycloalkyl)alkyl, phenylalkyl, alkoxy, alkylaminoalkyl, dialkylaminoalkyl, arylalkyl, aryloxyalkyl, arylalkyloxyalkyl, (heterocyclic radical)alkyl, and (heterocyclic radical)oxyalkyl. R<sub>1</sub> thus includes hydroxyalkyl, hydroxyalkenyl, hydroxyalkynyl, hydroxycycloalkyl, hydroxyaryl, aminoalkyl, aminoalkenyl, aminoalkynyl, aminocycloalkyl, aminoaryl, alkylalkenyl, (alkylaryl)alkyl, (haloaryl)alkyl, (hydroxyaryl)alkynyl, and so forth. Similarly, R<sub>A</sub> includes hydroxyalkyl and aminoaryl, and R<sub>B</sub> includes hydroxyalkyl, aminoalkyl, and hydroxyalkyl(heterocyclic radical)alkyl.

Heterocyclic radicals, which include but are not limited to heteroaryls, include: furyl, oxazolyl, isoxazolyl, thiophenyl, thiazolyl, pyrrolyl, imidazolyl, 1,3,4-triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyridazinyl, indolyl, and their nonaromatic counterparts. Further examples of heterocyclic radicals include piperidyl, quinolyl, isothiazolyl, piperidinyl, morpholinyl, piperazinyl, tetrahydrofuryl, tetrahydropyrrolyl, pyrrolidinyl, octahydroindolyl, octahydrobenzothiofuranyl, and octahydrobenzofuranyl.

Selective MEK 1 or MEK 2 inhibitors are those compounds which inhibit the MEK 1 or MEK 2 enzymes, respectively, without substantially inhibiting other enzymes such as MKK3, PKC, Cdk2A, phosphorylase kinase, EGF, and PDGF receptor kinases, and C-src. In general, a selective MEK 1 or MEK 2 inhibitor has an IC<sub>50</sub> for MEK 1 or MEK 2 that is at least one-fiftieth (1/50) that of its IC<sub>50</sub> for one of the above-named other enzymes. Preferably, a selective inhibitor has an IC<sub>50</sub> that is at least 1/100, more preferably 1/500, and even more preferably 1/1000, 1/5000, or less than that of its IC<sub>50</sub> or one or more of the above-named enzymes.

## B. Compounds

One aspect of the invention features the use of compounds shown in formula (I) in the Summary section. Embodiments of the invention includes compounds of formula (I) wherein: (a) Q is formula (i); (b) R<sub>3</sub> is H or fluoro; (c) R<sub>4</sub> is fluoro, chloro, or bromo; (d) R<sub>10</sub> is H, methyl, fluoro, or chloro; (e) R<sub>11</sub> is methyl, chloro, fluoro, nitro, or hydrogen; (f) R<sub>11</sub> is H; (g) R<sub>11</sub> is fluoro; (h) each of R<sub>10</sub> and R<sub>11</sub> is fluoro; (i) R<sub>1</sub> is H, methyl, ethyl, propyl, isopropyl, isobutyl, benzyl, phenethyl, allyl, C<sub>3-5</sub> alkenyl, C<sub>3-6</sub> cycloalkyl, (C<sub>3-5</sub> cycloalkyl)C<sub>1-2</sub> alkyl, (C<sub>3-5</sub> heterocyclic radical)C<sub>1-2</sub> alkyl, or (CH<sub>2</sub>)<sub>2-4</sub> NR<sub>C</sub>R<sub>D</sub>; (j) R<sub>1</sub> is H or (C<sub>3-4</sub> cycloalkyl)C<sub>1-2</sub> alkyl; (k) R<sub>2</sub> is H or methyl; (l) R<sub>A</sub> has at least one hydroxyl substituent; (m) R<sub>A</sub> is H, methyl, ethyl, isobutyl, hydroxyethyl, phenyl, 2-piperidin-1-yl-ethyl, 2,3-dihydroxy-propyl, 3-[4-(2-hydroxyethyl)-piperazin-1-yl]-propyl, 2-pyrrolidin-1-yl-ethyl, or 2-diethylamino-ethyl; and R<sub>B</sub> is H; or where R<sub>B</sub> is methyl and R<sub>A</sub> is phenyl.; (n) W is NR<sub>A</sub>R<sub>B</sub> or NR<sub>2</sub>NR<sub>A</sub>R<sub>B</sub>; (o) W is NR<sub>2</sub>(CH<sub>2</sub>)<sub>2-4</sub> NR<sub>A</sub>R<sub>B</sub> or O(CH<sub>2</sub>)<sub>2-3</sub> NR<sub>A</sub>R<sub>B</sub>; (p) W is NR<sub>2</sub>OR<sub>1</sub>; (q) W is OR<sub>1</sub>; (r) Z is formula (v); or (s) X<sub>1</sub> is NR<sub>8</sub>, and R<sub>7</sub> is H; or (t) combinations thereof. In formula (I), the values for Z are shown left to right, or in a counter-clockwise orientation around the phenyl ring of Q.

According to one aspect of the invention, the compound of formula (I) has a structure wherein: Q is formula (i) or (ii); R<sub>3</sub> is H or fluoro; R<sub>4</sub> is fluoro, chloro, or bromo; R<sub>10</sub> is H, methyl, or chloro; R<sub>11</sub> is chloro, fluoro, or hydrogen; R<sub>1</sub> is H, methyl, ethyl, propyl, isopropyl, isobutyl, benzyl, phenethyl, allyl, C<sub>3-5</sub> alkenyl, C<sub>3-6</sub> cycloalkyl, (C<sub>3-5</sub> cycloalkyl)C<sub>1-2</sub> alkyl, (C<sub>3-5</sub> heterocyclic radical)C<sub>1-2</sub> alkyl, or (CH<sub>2</sub>)<sub>2-4</sub> NR<sub>C</sub>R<sub>D</sub>; R<sub>1</sub> is H or (C<sub>3-4</sub> cycloalkyl)C<sub>1-2</sub> alkyl; R<sub>2</sub> is H or methyl; and Z is formula (v) or (vi). One embodiment of this aspect, X<sub>1</sub> is NR<sub>8</sub>. An example would be 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1[(2'-morpholinyl)-ethyl]-2-(phenyl)-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide.

Embodiments of the invention also include compounds wherein R<sub>10</sub> is H; R<sub>10</sub> is methyl or chloro; and where R<sub>10</sub> is chloro. In some embodiments, R<sub>7</sub> and R<sub>8</sub> together have no more than 14 carbon atoms, exclusive of R<sub>L</sub>, R<sub>M</sub>, R<sub>J</sub> and R<sub>K</sub>. Examples of this include compounds wherein R<sub>7</sub> and R<sub>8</sub> together

have no more than 13 carbon atoms; no more than 7, 8, or 10 carbon atoms; between 4 and 8 carbon atoms; between 1 and 10 carbon atoms; between 1 and 8 carbon atoms; and no more than 6 carbon atoms.

Preferably, where one of  $R_1$ ,  $R_2$ ,  $R_A$ ,  $R_B$ ,  $R_C$ ,  $R_D$ ,  $R_E$ ,  $R_F$ ,  $R_I$ ,  $R_J$ ,  $R_K$ ,  $R_L$ ,  $R_M$ ,  $R_G$ ,  $R_H$ ,  $R_N$ ,  $R_O$ , and  $R_P$  is an alkenyl or alkynyl group, its double or triple bond, respectively, is not adjacent the point of attachment. For example, where  $W$  is  $NR_2OR_1$ ,  $R_2$  is preferably prop-2-ynyl, or but-2 or 3-enyl, and less preferably prop-1-ynyl or but-1-enyl.

Listed below are some of the preferred structures which can be synthesized utilizing Schemes 1, 2, 10, and 11. Free acids, free hydroxamic acids, and cyclopropylmethyl hydroxamates are grouped together. For example, compounds 1, 11, and 21 differ only by "W" (as defined in the claims); compounds 2, 12, and 22 are similarly related. Preferred compounds also include the 2-chloro (replacing 2-methyl) analogs of the listed compounds.

Examples of compounds include: 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzimidazole-5-carboxylic acid (APK  $IC_{50} = 47 \pm 17$  nM); 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzooxazole-5-carboxylic acid; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzothiazole-5-carboxylic acid; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic acid; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylic acid; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-2-(2-hydroxyethyl)-1H-benzimidazole-5-carboxylic acid; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-2-(2-dimethylamino-ethyl)-1H-benzimidazole-5-carboxylic acid; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1-acetyl-benzimidazole-5-carboxylic acid; 8-Fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-carboxylic acid; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzimidazole-5-carboxylic acid hydroxyamide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzooxazole-5-carboxylic acid hydroxyamide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzothiazole-5-carboxylic acid hydroxyamide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic

acid hydroxyamide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylic acid hydroxyamide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-2-(2-hydroxyethyl)-1H-benzimidazole-5-carboxylic acid hydroxyamide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-2-(2-dimethylamino-ethyl)-1H-benzimidazole-5-carboxylic acid hydroxyamide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1-acetyl-benzimidazole-5-carboxylic acid hydroxyamide; 8-Fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid hydroxyamide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-carboxylic acid hydroxyamide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzooxazole-5-carboxylic acid cyclopropylmethoxy-amide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzothiazole-5-carboxylic acid cyclopropylmethoxy-amide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic acid cyclopropylmethoxy-amide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylic acid cyclopropylmethoxy-amide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-2-(2-hydroxyethyl)-1H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-2-(2-dimethylamino-ethyl)-1H-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide; 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1-acetyl-benzimidazole-5-carboxylic acid cyclopropylmethoxy-amide; 8-Fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid cyclopropylmethoxy-amide; and 7-Fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-carboxylic acid cyclopropylmethoxy-amide.

The following is a list of examples representing schemes 3-9. As above, free acids, free hydroxamic acids, and cyclopropylmethyl hydroxamates are grouped together. For example, compounds 31, 45, and 59 differ only by "W" (as defined in the claims); compounds 32, 46, and 60 are similarly related. Preferred compounds also include the 2-chloro (replacing 2-methyl) analogs of the listed compounds.

Examples of compounds from schemes 3-9 include: 4-Fluoro-5-(4-iodo-2-methyl-phenylamino)-benzothiazole-6-carboxylic acid; 4-Fluoro-5-(4-


iodo-2-methyl-phenylamino)-benzooxazole-6-carboxylic acid; 5-(2-Chloro-4-iodo-phenylamino)-6,7-difluoro-3H-benzoimidazole-4-carboxylic acid; 6,7-Difluoro-2-(2-hydroxy-ethyl)-5-(4-iodo-2-methyl-phenylamino)-3H-benzoimidazole-4-carboxylic acid; 6,7-Difluoro-5-(4-iodo-2-methyl-phenylamino)-benzooxazole-4-carboxylic acid; 6,7-Difluoro-5-(4-iodo-2-methyl-phenylamino)-benzothiazole-4-carboxylic acid; 7,8-Difluoro-6-(4-iodo-2-methyl-phenylamino)-quinoxaline-5-carboxylic acid; 6-(4-Iodo-2-methyl-phenylamino)-8-nitro-quinoxaline-5-carboxylic acid; 5-(4-Iodo-2-methyl-phenylamino)-8-nitro-quinoxaline-6-carboxylic acid; 8-Chloro-5-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid; 3-Cyclopropyl-7-(4-iodo-2-methyl-phenylamino)-3H-benzoimidazole-4,6-dicarboxylic acid 4-dimethylamide; 7-Bromo-4-(4-iodo-2-methyl-phenylamino)-benzooxazole-5-carboxylic acid; 7-(2-Chloro-4-iodo-phenylamino)-4-fluoro-benzothiazole-6-carboxylic acid; 7-(4-Iodo-2-methyl-phenylamino)-4-nitro-benzooxazole-6-carboxylic acid; 4-Fluoro-5-(4-iodo-2-methyl-phenylamino)-benzothiazole-6-carboxylic acid hydroxyamide; 4-Fluoro-5-(4-iodo-2-methyl-phenylamino)-benzooxazole-6-carboxylic acid hydroxyamide; 5-(2-Chloro-4-iodo-phenylamino)-6,7-difluoro-3H-benzoimidazole-4-carboxylic acid hydroxyamide; 6,7-Difluoro-2-(2-hydroxy-ethyl)-5-(4-iodo-2-methyl-phenylamino)-3H-benzoimidazole-4-carboxylic acid hydroxyamide; 6,7-Difluoro-5-(4-iodo-2-methyl-phenylamino)-benzooxazole-4-carboxylic acid hydroxyamide; 6,7-Difluoro-5-(4-iodo-2-methyl-phenylamino)-benzothiazole-4-carboxylic acid hydroxyamide; 7,8-Difluoro-6-(4-iodo-2-methyl-phenylamino)-quinoxaline-5-carboxylic acid hydroxyamide; 6-(4-Iodo-2-methyl-phenylamino)-8-nitro-quinoxaline-5-carboxylic acid hydroxyamide; 5-(4-Iodo-2-methyl-phenylamino)-8-nitro-quinoxaline-6-carboxylic acid hydroxyamide; 8-Chloro-5-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid hydroxyamide; 3-Cyclopropyl-7-(4-iodo-2-methyl-phenylamino)-3H-benzoimidazole-4,6-dicarboxylic acid 4-dimethylamide 6-hydroxyamide; 7-Bromo-4-(4-iodo-2-methyl-phenylamino)-benzooxazole-5-carboxylic acid hydroxyamide; 7-(2-Chloro-4-iodo-phenylamino)-4-fluoro-benzothiazole-6-carboxylic acid hydroxyamide; 7-(4-Iodo-2-methyl-phenylamino)-4-nitro-

benzooxazole-6-carboxylic acid hydroxyamide; 4-Fluoro-5-(4-iodo-2-methyl-phenylamino)-benzothiazole-6-carboxylic acid cyclopropylmethoxy-amide; 4-Fluoro-5-(4-iodo-2-methyl-phenylamino)-benzooxazole-6-carboxylic acid cyclopropylmethoxy-amide; 5-(2-Chloro-4-iodo-phenylamino)-6,7-difluoro-3H-benzoimidazole-4-carboxylic acid cyclopropylmethoxy-amide; 6,7-Difluoro-2-(2-hydroxy-ethyl)-5-(4-iodo-2-methyl-phenylamino)-3H-benzoimidazole-4-carboxylic acid cyclopropylmethoxy-amide; 6,7-Difluoro-5-(4-iodo-2-methyl-phenylamino)-benzooxazole-4-carboxylic acid cyclopropylmethoxy-amide; 6,7-Difluoro-5-(4-iodo-2-methyl-phenylamino)-benzothiazole-4-carboxylic acid cyclopropylmethoxy-amide; 7,8-Difluoro-6-(4-iodo-2-methyl-phenylamino)-quinoxaline-5-carboxylic acid cyclopropylmethoxy-amide; 6-(4-iodo-2-methyl-phenylamino)-8-nitro-quinoxaline-5-carboxylic acid cyclopropylmethoxy-amide; 5-(4-iodo-2-methyl-phenylamino)-8-nitro-quinoxaline-6-carboxylic acid cyclopropylmethoxy-amide; 8-Chloro-5-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid cyclopropylmethoxy-amide; 3-Cyclopropyl-7-(4-iodo-2-methyl-phenylamino)-3H-benzoimidazole-4,6-dicarboxylic acid 4-dimethylamide 6-cyclopropylmethoxy-amide; 7-Bromo-4-(4-iodo-2-methyl-phenylamino)-benzooxazole-5-carboxylic acid cyclopropylmethoxy-amide; 7-(2-Chloro-4-iodo-phenylamino)-4-fluoro-benzothiazole-6-carboxylic acid cyclopropylmethoxy-amide; and 7-(4-iodo-2-methyl-phenylamino)-4-nitro-benzooxazole-6-carboxylic acid cyclopropylmethoxy-amide.

## C. Synthesis


The disclosed compounds can be synthesized according to the following eleven Schemes, or variants thereof. These synthetic strategies are further exemplified in Examples 1-22 below.

Scheme 1







## Scheme 2




## Scheme 3




Scheme 4




## Scheme 5




## Scheme 6




Scheme 7



Scheme 8




Scheme 9






## Scheme 10



## Scheme 11



#### D. Uses

The disclosed compositions are useful as both prophylactic and therapeutic treatments for diseases or conditions relating to chronic pain, including neuropathic pain, as provided in the Summary section, as well as diseases or conditions modulated by the MEK cascade. For example, in one embodiment, the disclosed method relates to postoperative pain, phantom limb pain, burn pain, gout, trigeminal neuralgia, acute herpetic and postherpetic pain, causalgia, diabetic neuropathy, plexus avulsion, neuroma, vasculitis, crush injury, constriction injury, tissue injury, post-surgical pain, arthritis pain, or limb amputation

For example, local injuries can be treated with local or topical administration. Chronic pain affecting the entire body, such as diabetic neuropathy can be treated with systemic administration (injection or orally) of a disclosed composition. Treatment for chronic pain (e.g., post-operative pain) confined to the lower body can be administered centrally, e.g., epidurally. Formulations and methods of administration can include the use of more than one MEK inhibitor, or a combination of a MEK inhibitor and another pharmaceutical agent, such as an anti-inflammatory, analgesic, muscle relaxing, or anti-infective agent. Preferred routes of administration are oral, intrathecal or epidural, subcutaneous, intravenous, intramuscular, and, for non-human mammals, intraplantar, and are preferably epidural.

##### 1. Dosages

Those skilled in the art will be able to determine, according to known methods, the appropriate dosage for a patient, taking into account factors such as age, weight, general health, the type of pain requiring treatment, and the presence of other medications. In general, an effective amount will be between 0.1 and 1000 mg/kg per day, preferably between 1 and 300 mg/kg body weight, and daily dosages will be between 10 and 5000 mg for an adult subject of normal weight. Commercially available capsules or other

formulations (such as liquids and film-coated tablets) of 100 mg, 200 mg, 300 mg, or 400 mg can be administered according to the disclosed methods.

## 2. Formulations

Dosage unit forms include tablets, capsules, pills, powders, granules, aqueous and nonaqueous oral solutions and suspensions, and parenteral solutions packaged in containers adapted for subdivision into individual doses. Dosage unit forms can also be adapted for various methods of administration, including controlled release formulations, such as subcutaneous implants. Administration methods include oral, rectal, parenteral (intravenous, intramuscular, subcutaneous), intracisternal, intravaginal, intraperitoneal, intravesical, local (drops, powders, ointments, gels, or cream), and by inhalation (a buccal or nasal spray).

Parenteral formulations include pharmaceutically acceptable aqueous or nonaqueous solutions, dispersion, suspensions, emulsions, and sterile powders for the preparation thereof. Examples of carriers include water, ethanol, polyols (propylene glycol, polyethylene glycol), vegetable oils, and injectable organic esters such as ethyl oleate. Fluidity can be maintained by the use of a coating such as lecithin, a surfactant, or maintaining appropriate particle size. Carriers for solid dosage forms include (a) fillers or extenders, (b) binders, (c) humectants, (d) disintegrating agents, (e) solution retarders, (f) absorption accelerators, (g) adsorbants, (h) lubricants, (i) buffering agents, and (j) propellants.

Compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents; antimicrobial agents such as parabens, chlorobutanol, phenol, and sorbic acid; isotonic agents such as a sugar or sodium chloride; absorption-prolonging agents such as aluminum monostearate and gelatin; and absorption-enhancing agents.

## 3. Related compounds

The invention provides the disclosed compounds and closely related, pharmaceutically acceptable forms of the disclosed compounds, such as salts, esters, amides, hydrates or solvated forms thereof; masked or protected forms; and racemic mixtures, or enantiomerically or optically pure forms.

Pharmaceutically acceptable salts, esters, and amides include carboxylate salts (e.g., C<sub>1-8</sub> alkyl, cycloalkyl, aryl, heteroaryl, or non-aromatic heterocyclic), amino acid addition salts, esters, and amides which are within a reasonable benefit/risk ratio, pharmacologically effective, and suitable for contact with the tissues of patients without undue toxicity, irritation, or allergic response. Representative salts include hydrobromide, hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, valerate, oleate, palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulfonate. These may include alkali metal and alkali earth cations such as sodium, potassium, calcium, and magnesium, as well as non-toxic ammonium, quaternary ammonium, and amine cations such as tetramethyl ammonium, methylamine, trimethylamine, and ethylamine. See, for example, S.M. Berge, et al., "Pharmaceutical Salts," *J. Pharm. Sci.*, 1977, 66:1-19 which is incorporated herein by reference. Representative pharmaceutically acceptable amides of the invention include those derived from ammonia, primary C<sub>1-6</sub> alkyl amines and secondary di (C<sub>1-6</sub> alkyl) amines. Secondary amines include 5- or 6-membered heterocyclic or heteroaromatic ring moieties containing at least one nitrogen atom and optionally between 1 and 2 additional heteroatoms. Preferred amides are derived from ammonia, C<sub>1-3</sub> alkyl primary amines, and di (C<sub>1-2</sub> alkyl)amines. Representative pharmaceutically acceptable esters of the invention include C<sub>1-7</sub> alkyl, C<sub>5-7</sub> cycloalkyl, phenyl, and phenyl(C<sub>1-6</sub>)alkyl esters. Preferred esters include methyl esters.

The invention also includes disclosed compounds having one or more functional groups (e.g., hydroxyl, amino, or carboxyl) masked by a protecting group. Some of these masked or protected compounds are pharmaceutically acceptable; others will be useful as intermediates. Synthetic intermediates and processes disclosed herein, and minor modifications thereof, are also within the scope of the invention.

## HYDROXYL PROTECTING GROUPS

Hydroxyl protecting groups include: ethers, esters, and protection for 1,2- and 1,3-diols. The ether protecting groups include: methyl, substituted methyl ethers, substituted ethyl ethers, substituted benzyl ethers, silyl ethers and  
 5 conversion of silyl ethers to other functional groups.

### Substituted Methyl Ethers

Substituted methyl ethers include: methoxymethyl, methylthiomethyl, *t*-  
 utylthiomethyl, (phenyldimethylsilyl) methoxymethyl, benzyloxymethyl, *p*-  
 ethoxybenzyloxymethyl, (4-methoxyphenoxy) methyl, guaiacolmethyl, *t*-  
 10 butoxymethyl, 4-pentenylloxymethyl, siloxymethyl, 2-methoxyethoxymethyl,  
 2,2,2-trichloroethoxymethyl, bis(2-chloro- ethoxy)methyl, 2-  
 (trimethylsilyl)ethoxymethyl, tetrahydropyranyl, 3-bromotetrahydro-pyranyl,  
 tetrahydrothiopyranyl, 1-methoxycyclohexyl, 4-methoxytetrahydropyranyl, 4-  
 methoxytetrahydrothio-pyranyl, 4-methoxytetrahydrothiopyranyl *S,S*-dioxido,  
 15 1-[(2-chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl, 1,4-dioxan-2-yl,  
 tetrahydrofuranyl, tetrahydrothiofuranyl, and 2,3,3a,4,5,6,7,7a-octahydro-  
 7,8,8-trimethyl-4,7-ethanobenzofuran-2-yl.

### Substituted Ethyl Ethers

Substituted ethyl ethers include: 1-ethoxyethyl, 1-(2, chloroethoxy)ethyl,  
 20 1-methyl-1-methoxyethyl, 1-methyl-1-benzyloxyethyl, 1-methyl-1-benzyloxy-2-  
 fluoroethyl, 2,2,2-trichloroethyl, 2-trimethylsilyl ethyl, 2-(phenylselenyl)ethyl, *t*-  
 butyl, allyl, *p*-chlorophenyl, *p*-methoxyphenyl, 2,4-dinitrophenyl, and benzyl.

### Substituted Benzyl Ethers

Substituted benzyl ethers include: *p*-methoxybenzyl, 3,4-dimethoxybenzyl,  
 25 *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-halobenzyl, 2,6-dichlorobenzyl, *p*-cyanobenzyl,  
*p*-phenylbenzyl, 2- and 4-picolyl, 3-methyl-2-picolyl *N*-oxido, diphenylmethyl,  
*p*, *p'*-dinitrobenzhydryl, 5-dibenzosuberyl, triphenylmethyl,  $\alpha$ -naphthyldiphenyl-  
 methyl, *p*-methoxyphenyldiphenylmethyl, di(*p*-methoxyphenyl)phenylmethyl,  
 tri-(*p*-methoxyphenyl)methyl, 4-(4'-bromophenacyloxy)phenyldiphenylmethyl,  
 30 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-  
 tris(levulinoyloxyphenyl) methyl, 4,4',4''tris(benzoyloxyphenyl)methyl, 3-  
 (imidazol-1-ylmethyl)bis(4',4''-dimethoxyphenyl)-methyl, 1,1-bis(4-

methoxyphenyl)-1'-pyrenylmethyl, 9-anthryl, 9-(9-phenyl) xanthenyl, 9-(9-phenyl-10-oxo) anthryl, 1,3-benzodithiolan-2-yl, and benzisothiazolyl S,S-dioxido.

#### Silyl Ethers

5 Silyl ethers include: trimethylsilyl, triethylsilyl, triisopropylsilyl, dimethylisopropylsilyl, diethylisopropylsilyl, dimethylhexylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl, tribenzylsilyl, tri-*p*-xylylsilyl, triphenylsilyl, diphenylmethylsilyl, and *t*-butylmethoxyphenylsilyl.

#### 10 ESTERS

Esters protecting groups include: esters, carbonates, assisted cleavage, miscellaneous esters, and sulfonates.

#### Esters

15 Examples of protective esters include: formate, benzoylformate, acetate, chloroacetate, dichloroacetate, trichloroacetate, trifluoroacetate, methoxyacetate, triphenylmethoxyacetate, phenoxyacetate, *p*-chlorophenoxyacetate, *p*-P-phenylacetate, 3-phenylpropionate, 4-oxopentanoate (levulinate), 4,4-(ethylenedithio) pentanoate, pivaloate, adamantooate, crotonate, 4-methoxycrotonate, benzoate, *p*-phenylbenzoate, 20 and 2,4,6-trimethylbenzoate (mesitoate).

#### Carbonates

Carbonates include: methyl, 9-fluorenylmethyl, ethyl, 2,2,2-trichloroethyl, 2-(trimethylsilyl) ethyl, 2-(phenylsulfonyl) ethyl, 2-(triphenylphosphonio) ethyl, isobutyl, vinyl, allyl, *p*-nitrophenyl, benzyl, *p*-methoxybenzyl, 3,4- 25 dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, S-benzyl thiocarbonate, 4-ethoxy-1-naphthyl, and methyl dithiocarbonate.

#### Assisted Cleavage

30 Examples of assisted cleavage protecting groups include: 2-iodobenzoate, 4-azido-butyrates, 4-nitro-4-methylpentanoate, *o*-(dibromomethyl) benzoate, 2-formylbenzene-sulfonate, 2-(methylthiomethoxy) ethyl carbonate, 4-(methylthiomethoxymethyl) benzoate, and 2-(methylthiomethoxymethyl) benzoate.

### Miscellaneous Esters

In addition to the above classes, miscellaneous esters include: 2,6-dichloro-4-methylphenoxyacetate, 2,6-dichloro-4-(1,1,3,3-tetramethylbutyl) phenoxyacetate, 2,4-bis(1,1-dimethylpropyl) phenoxyacetate, 5 chlorodiphenylacetate, isobutyrate, monosuccinoate, (*E*)-2-methyl-2-butenate (tigloate), *o*-(methoxycarbonyl) benzoate, *p*-P-benzoate,  $\alpha$ -naphthoate, nitrate, alkyl *N,N,N',N'*-tetramethylphosphorodiamidate, *N*-phenylcarbamate, borate, dimethylphosphinothioyl, and 2,4-dinitrophenylsulfenate.

### Sulfonates

Protective sulfates includes: sulfate, methanesulfonate(mesylate), benzylsulfonate, and tosylate.

### PROTECTION FOR 1,2- AND 1,3-DIOLS

The protection for 1,2 and 1,3-diols group includes: cyclic acetals and ketals, cyclic ortho esters, and silyl derivatives.

### Cyclic Acetals and Ketals

Cyclic acetals and ketals include: methylene, ethylidene, 1-*t*-butylethylidene, 1-phenylethylidene, (4-methoxyphenyl) ethylidene, 2,2,2-trichloroethylidene, 20 acetonide (isopropylidene), cyclopentylidene, cyclohexylidene, cycloheptylidene, benzylidene, *p*-methoxybenzylidene, 2,4-dimethoxybenzylidene, 3,4-dimethoxybenzylidene, and 2-nitrobenzylidene.

### Cyclic Ortho Esters

Cyclic ortho esters include: methoxymethylene, ethoxymethylene, dimethoxy- 25 methylene, 1-methoxyethylidene, 1-ethoxyethylidene, 1,2-dimethoxyethylidene,  $\alpha$ -methoxybenzylidene, 1-(*N,N*-dimethylamino)ethylidene derivative,  $\alpha$ -(*N,N*-dimethylamino) benzylidene derivative, and 2-oxacyclopentylidene.



## PROTECTION FOR THE CARBOXYL GROUP

### ESTERS

Ester protecting groups include: esters, substituted methyl esters, 2-substituted ethyl esters, substituted benzyl esters, silyl esters, activated  
5 esters, miscellaneous derivatives, and stannyl esters.

#### Substituted Methyl Esters

Substituted methyl esters include: 9-fluorenylmethyl, methoxymethyl, methylthiomethyl, tetrahydropyranyl, tetrahydrofuranyl, methoxyethoxymethyl, 2-(trimethylsilyl)ethoxy-methyl, benzyloxymethyl, phenacyl, *p*-bromophenacyl,  
10  $\alpha$ -methylphenacyl, *p*-methoxyphenacyl, carboxamidomethyl, and *N*-phthalimidomethyl.

#### 2-Substituted Ethyl Esters

2-Substituted ethyl esters include: 2,2,2-trichloroethyl, 2-haloethyl,  $\alpha$ -chloroalkyl, 2-(trimethylsilyl)ethyl, 2-methylthioethyl, 1,3-dithianyl-2-methyl,  
15 2(*p*-nitrophenylsulfenyl)-ethyl, 2-(*p*-toluenesulfonyl)ethyl, 2-(2'-pyridyl)ethyl, 2-(diphenylphosphino)ethyl, 1-methyl-1-phenylethyl, *t*-butyl, cyclopentyl, cyclohexyl, allyl, 3-buten-1-yl, 4-(trimethylsilyl)-2-buten-1-yl, cinnamyl,  $\alpha$ -methylcinnamyl, phenyl, *p*-(methylmercapto)-phenyl, and benzyl.

#### Substituted Benzyl Esters

Substituted benzyl esters include: triphenylmethyl, diphenylmethyl, bis(*o*-nitrophenyl)methyl, 9-anthrylmethyl, 2-(9,10-dioxo)anthrylmethyl, 5-dibenzo-suberyl, 1-pyrenylmethyl, 2-(trifluoromethyl)-6-chromylmethyl, 2,4,6-trimethylbenzyl, *p*-bromobenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-methoxybenzyl, 2,6-dimethoxybenzyl, 4-(methylsulfinyl)benzyl, 4-sulfobenzyl,  
25 piperonyl, and 4-P-benzyl.

#### Silyl Esters

Silyl esters include: trimethylsilyl, triethylsilyl, *t*-butyldimethylsilyl, *i*-propyldimethylsilyl, phenyldimethylsilyl, and di-*t*-butylmethylsilyl.

#### Miscellaneous Derivatives

Miscellaneous derivatives includes: oxazoles, 2-alkyl-1,3-oxazolines, 4-alkyl-5-oxo-1,3-oxazolidines, 5-alkyl-4-oxo-1,3-dioxolanes, ortho esters, phenyl  
30 group, and pentaaminocobalt(III) complex.

### Stannyl Esters

Examples of stannyl esters include: triethylstannyl and tri-*n*-butylstannyl.

### AMIDES AND HYDRAZIDES

- 5 Amides include: *N,N*-dimethyl, pyrrolidinyl, piperidinyl, 5,6-dihydrophenanthridinyl, *o*-nitroanilides, *N*-7-nitroindolyl, *N*-8-nitro-1,2,3,4-tetrahydroquinolyl, and *p*-P-benzenesulfonamides. Hydrazides include: *N*-phenyl, *N,N'*-diisopropyl and other dialkyl hydrazides.

### 10 PROTECTION FOR THE AMINO GROUP

#### CARBAMATES

Carbamates include: carbamates, substituted ethyl, assisted cleavage, photolytic cleavage, urea-type derivatives, and miscellaneous carbamates.

#### 15 Carbamates

Carbamates include: methyl and ethyl, 9-fluorenylmethyl, 9-(2-sulfo)fluorenylmethyl, 9-(2,7-dibromo)fluorenylmethyl, 2,7-di-*t*-butyl-[9-(10,10-dioxo-10,10,10,10-tetrahydro- thioxanthyl)]methyl, and 4-methoxyphenacyl.

#### Substituted Ethyl

- 20 Substituted ethyl protective groups include: 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-phenylethyl, 1-(1-adamantyl)-1-methylethyl, 1,1-dimethyl-2-haloethyl, 1,1dimethyl-2,2-dibromoethyl, 1,1-dimethyl-2,2,2-trichloroethyl, 1-methyl-1-(4-biphenyl)ethyl, 1-(3,5-di-*t*-butylphenyl)-1-methylethyl, 2-(2'-and 4'-pyridyl)ethyl, 2-(*N,N*-icyclohexylcarboxamido)- ethyl, *t*-butyl, 1-adamantyl, 25 vinyl, allyl, 1-isopropylallyl, connamyl, 4-nitrocinnamyl, quinolyl, *N*-hydroxypiperidinyl, alkylidithio, benzyl, *p*-methoxybenzyl, *p*-nitrobenzyl, *p*-bromobenzyl, *p*-chlorobenzyl, 2,4dichlorobenzyl, 4-methylsulfinylbenzyl, 9-anthrylmethyl, and diphenylmethyl.

#### 30 Assisted Cleavage

Protection via assisted cleavage includes: 2-methylthioethyl, 2-methylsulfonylethyl, 2-(*p*-toluenesulfonyl)ethyl, [2-(1,3-dithianyl)]methyl,

4-methylthiophenyl, 2,4-dimethyl-thiophenyl, 2-phosphonioethyl, 2-triphenylphosphonioisopropyl, 1,1-dimethyl-2cyanoethyl, *m*-chloro-*p*-acyloxybenzyl, *p*-(dihydroxyboryl)benzyl, 5-benzisoxazolyl-methyl, and 2-(trifluoromethyl)-6-chromonylmethyl.

#### 5      Photolytic Cleavage

Photolytic cleavage methods use groups such as: *m*-nitrophenyl, 3,5-dimethoxybenzyl, *o*-nitrobenzyl, 3,4-dimethoxy-6-nitrobenzyl, and phenyl(*o*-nitrophenyl)methyl.

#### Urea-Type Derivatives

10      Examples of of urea-type derivatives include: phenothiazinyl-(10)-carbonyl derivative, *N*'-*p*-toluenesulfonylaminocarbonyl, and *N*'-phenylaminothiocarbonyl.

#### Miscellaneous Carbamates

15      In addition to the above, miscellaneous carbamates include: *t*-amyl, *S*-benzyl thiocarbamate, *p*-cyanobenzyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopropylmethyl, *p*-decyloxy-benzyl, diisopropylmethyl, 2,2-dimethoxycarbonylvinyl, *o*-(*N,N*-dimethyl-carboxamido)-benzyl, 1,1-dimethyl-3(*N,N*-dimethylcarboxamido)propyl, 1,1-dimethyl-propynyl, di(2-pyridyl)methyl, 2-furanylmethyl, 2-iodoethyl, isobornyl, isobutyl, isonicotinyl, *p*(*p*'-methoxyphenyl- azo)benzyl, 1-methylcyclobutyl, 1-methylcyclohexyl, 1-20 methyl-1-cyclopropyl- methyl, 1-methyl-(3,5-dimethoxyphenyl)ethyl, 1-methyl-1(*p*-henylazophenyl)- ethyl, 1-methyl-1-phenylethyl, 1-methyl-1-(4-pyridyl)ethyl, phenyl, *p*-(phenylazo)benzyl, 2,4,6-tri-*t*-butylphenyl, 4-(trimethylammonium) benzyl, and 2,4,6-trimethylbenzyl.

25

## AMIDES

### Amides

Amides includes: *N*-formyl, *N*-acetyl, *N*-chloroacetyl, *N*-trichloroacetyl, *N*-trifluoroacetyl, *N*-phenylacetyl, *N*-3-phenylpropionyl, *N*-picolinoyl, *N*-3-pyridyl-carboxamide, *N*-benzoylphenylalanyl derivative, *N*-benzoyl, and *N*-*p*-phenylbenzoyl.

### Assisted Cleavage

Assisted cleavage groups include: *N*-*o*-nitrophenylacetyl, *N*-*o*-nitrophenoxyacetyl, *N*-acetoacetyl, (*N*'-dithiobenzyloxycarbonylamino)acetyl, *N*-3-(*p*-hydroxyphenyl) propionyl, *N*-3-(*o*-nitrophenyl)propionyl, *N*-2-methyl-2-(*o*-nitrophenoxy)propionyl, *N*-2-methyl-2-(*o*-phenylazophenoxy)propionyl, *N*-4-chlorobutyryl, *N*-3-methyl-3-nitrobutyryl, *N*-*o*-nitrocinnamoyl, *N*-acetylmethionine derivative, *N*-*o*-nitrobenzoyl, *N*-*o*-(benzoyloxymethyl)benzoyl, and 4,5-diphenyl-3-oxazolin-2-one.

### Cyclic Imide Derivatives

Cyclic imide derivatives include: *N*-phthalimide, *N*-dithiasuccinoyl, *N*-2,3-diphenyl-maleoyl, *N*-2,5-dimethylpyrrolyl, *N*-1,1,4,4-tetramethyldisilylazacyclopentane adduct, 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, and 1-substituted 3,5-dinitro-4-pyridonyl.

## SPECIAL -NH PROTECTIVE GROUPS

Protective groups for –NH include: *N*-alkyl and *N*-aryl amines, imine derivatives, enamine derivatives, and *N*-hetero atom derivatives (such as *N*-metal, *N*-N, *N*-P, *N*-Si, and *N*-S), *N*-sulfenyl, and *N*-sulfonyl.

### *N*-Alkyl and *N*-Aryl Amines

*N*-alkyl and *N*-aryl amines include: *N*-methyl, *N*-allyl, *N*-[2-(trimethylsilyl)ethoxyl]-methyl, *N*-3-acetoxypentyl, *N*-(1-isopropyl-4-nitro-2-oxo-3-pyrrolin-3-yl), quaternary ammonium salts, *N*-benzyl, *N*-di(4-methoxyphenyl)methyl, *N*-5-dibenzosuberonyl, *N*-triphenylmethyl, *N*-(4-methoxyphenyl)diphenylmethyl, *N*-9-phenylfluorenyl,

*N*-2,7-dichloro-9-fluorenylmethylene, *N*-ferrocenylmethyl, and *N*-2-picolylamine *N*'-oxide.

#### Imine Derivatives

Imine derivatives include: *N*-1,1-dimethylthiomethylene, *N*-benzylidene,

5 *N*-*p*-methoxybenzylidene, *N*-diphenylmethylene,

*N*-[(2-pyridyl)mesityl]methylene,

*N*-(*N*',*N*'-dimethylaminomethylene), *N,N*'-isopropylidene,

*N*-*p*-nitrobenzylidene,

*N*-salicylidene, *N*-5-chlorosalicylidene, *N*-(5-chloro-2-hydroxyphenyl)phenyl-

10 methylene, and *N*-cyclohexylidene.

#### Enamine Derivative

An example of an enamine derivative is *N*-

(5,5-dimethyl-3-oxo-1-cyclohexenyl).

#### *N*-Hetero Atom Derivatives

15 *N*-metal derivatives include: *N*-borane derivatives, *N*-diphenylborinic acid derivative, *N*-[phenyl(pentacarbonylchromium- or -tungsten)]carbenyl, and

*N*-copper or *N*-zinc chelate. Examples of *N-N* derivatives include: *N*-nitro, *N*-nitroso, and *N*-oxide. Examples of *N-P* derivatives include:

*N*-diphenylphosphinyl, *N*-dimethylthiophosphinyl, *N*-diphenylthiophosphinyl,

20 *N*-dialkyl phosphoryl, *N*-dibenzyl phosphoryl, and *N*-diphenyl phosphoryl.

Examples of *N*-sulfenyl derivatives include: *N*-benzenesulfenyl,

*N*-*o*-nitrobenzenesulfenyl, *N*-2,4-dinitrobenzenesulfenyl,

*N*-pentachlorobenzenesulfenyl, *N*-2-nitro-4-methoxy-benzenesulfenyl,

*N*-triphenylmethylsulfenyl, and *N*-3-nitropyridinesulfenyl. *N*-sulfonyl

25 derivatives include: *N*-*p*-toluenesulfonyl, *N*-benzenesulfonyl,

*N*-2,3,6-trimethyl-4-methoxybenzenesulfonyl,

*N*-2,4,6-trimethoxybenzenesulfonyl, *N*-2,6-dimethyl-4-methoxy-

benzenesulfonyl, *N*-pentamethylbenzenesulfonyl,

*N*-2,3,5,6-tetramethyl-4-methoxybenzene- sulfonyl,

30 *N*-4-methoxybenzenesulfonyl, *N*-2,4,6-trimethylbenzenesulfonyl, *N*-

2,6-dimethoxy-4-methylbenzenesulfonyl,

*N*-2,2,5,7,8-pentamethylchroman-6-sulfonyl, *N*-methanesulfonyl,

*N*- $\beta$ -trimethylsilylethanesulfonyl, *N*-9-anthracenesulfonyl, *N*-4-(4',8'-dimethoxynaphthylmethyl)-benzenesulfonyl, *N*-benzylsulfonyl, *N*-trifluoromethylsulfonyl, and *N*-phenacylsulfonyl.

5            Disclosed compounds which are masked or protected may be prodrugs, compounds metabolized or otherwise transformed *in vivo* to yield a disclosed compound, e.g., transiently during metabolism. This transformation may be a hydrolysis or oxidation which results from contact with a bodily fluid such as blood, or the action of acids, or liver, gastrointestinal, or other enzymes.

10           Features of the invention are further described in the examples below.

## E. Examples

## BIOLOGICAL EXAMPLES

5

## Example 1

**Effect of PD 198306 on streptozocin-induced static allodynia**Animals

Male Sprague Dawley rats (250-300g), obtained from Bantin and Kingman, (Hull, U.K.) were housed in groups of 3. All animals were kept under a 12h light/dark cycle (lights on at 07h 00min) with food and water *ad libitum*. All experiments were carried out by an observer blind to drug treatments.

Development of diabetes in the rat

Diabetes was induced in rats by a single i.p. injection of streptozocin (50 mg/kg) as described previously (Courteix et al., 1993).

Evaluation of static allodynia

Mechanical hypersensitivity was measured using Semmes-Weinstein von Frey hairs (Stoelting, Illinois, U.S.A.). Animals were placed into wire mesh bottom cages allowing access to the underside of their paws. Animals were habituated to this environment prior to the start of the experiment. Mechanical hypersensitivity was tested by touching the plantar surface of the animals right hind paw with von Frey hairs in ascending order of force ( 0.7, 1.2, 1.5, 2, 3.6, 5.5, 8.5, 11.8, 15.1 and 29g) for up to 6 sec. Once a withdrawal response was established, the paw was re-tested, starting with the next descending von Frey hair until no response occurred. The highest force of 29 g lifted the paw as well as eliciting a response, thus represented the cut off point. The lowest amount of force required to elicit a response was recorded as the paw withdrawal threshold (PWT) in grams.

### Drugs

PD 198306 [N-Cyclopropylmethoxy-3,4,5-trifluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide] and CI-1008 (pregabalin) were synthesized at Parke-Davis (Ann Arbor, MI, USA). PD 198306 was suspended in cremophor:ethanol:water (1:1:8) vehicle. Pregabalin was dissolved in water. Both compounds were administered orally. Streptozocin (Aldrich, UK) was dissolved in 0.9% w/v NaCl and administered intraperitoneally. Drug administrations were made in a volume of 1 ml/kg.

### Statistics

The static allodynia data were analysed using a Kruskal-Wallis ANOVA for non-parametric results, followed when significant by Mann-Whitney's t test.

### Experimental protocol

Static allodynia was assessed with von Frey hairs, before (baseline, BL) and 1h after oral administration of PD 198306 (30mg/kg, p.o.), vehicle (cremophor:ethanol:water, 1:1:8) or pregabalin (30mg/kg, p.o.) (test). Animals were administered again the same compounds on the following day, both in the morning and the afternoon. Static allodynia was assessed only before and 1h after the afternoon administration, in order to minimise the habituation of the animals to the testing conditions. Animals treated with pregabalin received water in the morning administration, in order to avoid the potential development of tolerance to the compound with repeated administration.

#### Day 1:

p.m.: BL

#### Day 2:

a.m.: PD 198306

Water

Vehicle

p.m.: BL



PD 198306

PD 198306

Pregabalin

Pregabalin

Vehicle

Vehicle

**Test****Test**

5

**RESULTS**

A single administration of pregabalin (30mg/kg, p.o.) significantly blocked streptozocin-induced static allodynia 1h after administration. In contrast, a single administration of PD 198306 (30mg/kg, p.o) had no effect on streptozocin-induced static allodynia 1h after administration (see below). However, after the compound had been administered twice more on the following day, it significantly blocked streptozocin-induced static allodynia 1h after the third administration. The effects had disappeared by the following day (see FIG. 1).

15

**Example 2****MATERIALS AND METHODS****Animals**

Male Sprague Dawley rats (250-300g), obtained from Charles River, Margate, U.K.) were housed in groups of 3-6. All animals were kept under a 12h light/dark cycle (lights on at 07h 00min) with food and water *ad libitum*. All experiments were carried out by an observer blind to drug treatments.

Diabetes was induced in rats by a single i.p. injection of streptozocin (50mg/kg) as described previously (Courteix et al., 1993).

25

**Development of Chronic Constriction Injury in the rat**

Animals were anaesthetised with 2% isoflurane 1:4 O<sub>2</sub>/N<sub>2</sub>O mixture maintained during surgery via a nose cone. The sciatic nerve was ligated as previously described by Bennett and Xie, 1988. Animals were placed on a homeothermic blanket for the duration of the procedure. After surgical preparation the common sciatic nerve was exposed at the middle of the thigh

30

by blunt dissection through biceps femoris. Proximal to the sciatic trifurcation, about 7mm of nerve was freed of adhering tissue and 4 ligatures (4-0 silk) were tied loosely around it with about 1mm spacing. The incision was closed in layers and the wound treated with topical antibiotics.

5

#### Intrathecal injections

PD 198306 and pregabalin were administered intrathecally in a volume of 10  $\mu$ l using a 100  $\mu$ l Hamilton syringe by exposing the spine of the rats under brief isoflurane anaesthesia. Injections were made into the intrathecal space between lumbar region 5-6 with a 10 mm long 27 gauge needle. Penetrations were judged successful if there was a tail flick response. The wound was sealed with an autoclip and rats appeared fully awake within 2-3 min following injection.

10

#### Evaluation of static allodynia

Mechanical hypersensitivity was measured using Semmes-Weinstein von Frey hairs (Stoelting, Illinois, U.S.A.). Animals were placed into wire mesh bottom cages allowing access to the underside of their paws. Animals were habituated to this environment prior to the start of the experiment. Mechanical hypersensitivity was tested by touching the plantar surface of the animals right hind paw with von Frey hairs in ascending order of force ( 0.7, 1.2, 1.5, 2, 3.6, 5.5, 8.5, 11.8, 15.1 and 29g) for up to 6sec. Once a withdrawal response was established, the paw was re-tested, starting with the next descending von Frey hair until no response occurred. The highest force of 29g lifted the paw as well as eliciting a response, thus represented the cut off point. The lowest amount of force required to elicit a response was recorded as the paw withdrawal threshold (PWT) in grams.

15

20

25

#### Experimental protocol

Static allodynia was assessed with von Frey hairs, before (baseline, BL) and 0.5h, 1h and 2h after intrathecal or intraplantar administration of PD 198306 (1-30 $\mu$ g, i.t.), vehicle (cremophor:ethanol:water, 1:1:8) or pregabalin (10 $\mu$ g, i.t). For oral administration experiments, static allodynia was assessed

30

with von Frey hairs, before (baseline, BL) and 1h after oral administration of PD 198306 (3-30mg/kg, p.o.), vehicle (cremophor:ethanol:water, 1:1:8) or pregabalin (30mg/kg, p.o.). Animals were administered again the same compounds on the following day, both in the morning and the afternoon. Static allodynia was assessed before and 1h after the morning administration. In the afternoon static allodynia was assessed before, 1h, 2h and 3h after administration for streptozocin treated animals. CCI animals were assessed before, 1h and 2h after administration

#### 10 Drugs used

PD 198306 and pregabalin were synthesised at Parke-Davis (Ann Arbor, MI, USA). PD 198306 was suspended in cremophor:ethanol:water (1:1:8) vehicle. Pregabalin was dissolved in water. Both compounds were administered orally, intrathecally or intraplantar in volumes of 1ml/kg, 10µl and 15 100µl respectively. Streptozocin (Aldrich, UK) was dissolved in 0.9% w/v NaCl and administered intraperitoneally in a volume of 1ml/kg.

#### Statistics

Data were analysed using a Kruskal-Wallis ANOVA for non-parametric results, followed when significant by Mann-Whitney's t test vs vehicle group.

### RESULTS

#### **1. Effects of PD 198306 on static allodynia, following systemic administration**

##### 25 1.1. Effect of PD198306 on streptozocin-induced static allodynia

A single administration of pregabalin (30mg/kg, p.o.) significantly blocked streptozocin-induced static allodynia 1h after administration. In contrast, a single administration of PD 198306 (3-30mg/kg, p.o) had no effect on streptozocin-induced static allodynia 1h after administration (FIG. 2).

30 However, after the compound had been administered twice more on the following day, PD 198306 (30mg/kg) significantly blocked streptozocin-induced static allodynia for 2h after the third administration (FIG. 2).

### 1.2. Effect of PD198306 on CCI-induced static allodynia

A single administration of pregabalin (30mg/kg, p.o.) significantly blocked CCI-induced static allodynia 1h after administration. In contrast, neither a single or  
5 multiple administration of PD 198306 (3-30mg/kg, p.o) had any effect on CCI-induced static allodynia (FIG. 3).

### **2. Effects of PD 198306 on static allodynia, following intrathecal administration**

10 Intrathecally administered PD198306 (1-30 $\mu$ g) dose-dependently blocked the maintenance of static allodynia in both streptozocin (FIG. 4) and CCI animals (FIG. 5) with respective MEDs of 3 and 10  $\mu$ g. This antiallodynic effect lasted for 1h.

### **3. Effects of PD 198306 on static allodynia, following intraplantar administration**

15 An intrathecal administration of PD 198306 (30 $\mu$ g) significantly blocked static allodynia in both neuropathic pain models (FIGS. 6,7). In contrast, a single administration of PD 198306 at a dose 100-fold higher (3mg/100 $\mu$ l) directly  
20 into the paw had no effect on streptozocin (FIG. 6) or CCI-induced static allodynia (FIG. 7).

### REFERENCES

Bennett GJ, Xie Y-K. A peripheral mononeuropathy in rat that produces  
25 disorders of pain sensation like those seen in man. Pain 1988;33:87-107.

Courteix C, Eschalier A and Lavarenne J. Streptozocin –induced rats: behavioural evidence for a model of chronic pain. Pain 1993;53:81-8

## Example 3

**Effect of other MEK inhibitors in a neuropathic pain model in the rat**SUMMARY

5       The effect of several MEK inhibitors, with different binding affinities, has been investigated in the CCI model of neuropathic pain in the rat, by assessing static allodynia with von Frey hairs. Intrathecal administration of PD219622 or PD297447 (30µg) had no significant effect on allodynia. This lack of effect may reflect the low affinity or solubility of the compounds. However, 10 intrathecal administration of PD 254552 or PD 184352 (30µg), which possesses higher binding affinities, blocked the maintenance of static allodynia in CCI animals. The antiallodynic effect was only evident for 30min post-injection and thus, shorter than the one observed for pregabalin (100µg). The magnitude of the effect was similar for 30µg of PD 184352 and 100µg of pregabalin. From 15 this study it is concluded that MEK inhibitors exert an antiallodynic effect in CCI-induced neuropathic rats when administered intrathecally, and that the antiallodynic effect correlates with the affinity of the compounds.

20       The animals and methods for developing chronic constriction injury in the rat, injecting test compounds, and evaluation of static allodynia were according to Example 2 above. PD219622, PD297447, PD 184352, PD 254552 and pregabalin were administered intrathecally at doses of 30µg for all PD compounds and 100µg for pregabalin. Static allodynia was assessed with von Frey hairs, before (baseline, BL) and 0.5h, 1h and 2h after intrathecal 25 administration of the compounds

Drugs used

PD297447, PD219622, PD 254552, PD 184352 (CI-1040), and pregabalin were synthesised at Parke-Davis (Ann Arbor, MI, USA). PD297447, 30 PD219622, PD 254552 and PD 184352 were suspended in cremophor:ethanol:water (1:1:8) vehicle. Pregabalin was dissolved in water. All compounds were administered intrathecally in a 10µl volume.

### Statistics

Data were analysed using a Kruskal-Wallis ANOVA for non-parametric results, followed when significant by Mann-Whitney's t test vs vehicle group.

5

### RESULTS

10

Intrathecaly administered PD297447 or PD219622 (30 $\mu$ g) had no significant effect on allodynia. This lack of effect may reflect the low affinity of the compounds (965nM and 100nM respectively). However, intrathecal administration of PD 184352 or PD 254552 (30 $\mu$ g) blocked the maintenance of static allodynia in CCI animals (see FIG. 8). These compounds possess higher affinity (2 and 5 nM respectively). The antiallodynic effect was only evident for 30min post-injection and thus, shorter than the one observed for pregabalin (100 $\mu$ g). The magnitude of the effect was similar for 30 $\mu$ g of PD 184352 and 100 $\mu$ g of pregabalin.

15

The results indicate that MEK inhibitors exert an antiallodynic effect in CCI-induced neuropathic rats when administered intrathecaly, and that the antiallodynic effect correlates with the affinity of the compounds.

CHEMICAL EXAMPLES

## EXAMPLE 1

- 5 Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzimidazole-5-carboxylic acid (PD 205293) (APK IC<sub>50</sub> = 14 nM; colon 26 cells, IC<sub>50</sub> = > 10 micromolar)

Step a: Preparation of 5-nitro-2,3,4-trifluorobenzoic acid

- 10 To gently stirring concentrated sulfuric acid (50 ml) was added fuming nitric acid (3.4 ml, 0.076 mol). Solid 2,3,4-trifluorobenzoic acid (10.00 g, 0.05565 mol) was added directly in increments. After stirring 45 minutes, the reaction mixture had become an orange homogeneous solution which was then poured over chilled water (400 ml). The resulting aqueous suspension
- 15 was extracted with diethyl ether (3 x 200 ml). The combined extracts were dried with anhydrous magnesium sulfate and concentrated *in vacuo* to yield 12.30 g of a dull, light-yellow solid. Recrystallization from chloroform (50 ml) afforded 9.54 g of the pale yellow microcrystalline product; 78 % yield; m.p. ;
- 20 <sup>1</sup>H-NMR (400 MHz; DMSO) δ 14.29 (broad s, 1H), 8.43-8.38 (m, 1H); <sup>13</sup>C-NMR (100 MHz; DMSO) δ 162.41, 154.24 (dd, J<sub>C-F</sub>=270.1, 10.7 Hz), 148.35 (dd, J<sub>C-F</sub>=267.0, 9.2 Hz), 141.23 (dt, J<sub>C-F</sub>=253.4 Hz), 133.95, 123.30 (d, J<sub>C-F</sub>=2.2 Hz), 116.92 (dd, J<sub>C-F</sub>=18.2, 3.8 Hz); <sup>19</sup>F-NMR (376 MHz; DMSO) δ -120.50 to -120.63 (m), -131.133 to -131.27 (m), -153.63 to -153.74 (m).

- 25 Step b: Preparation of 4-amino-2,3-difluoro-5-nitrobenzoic acid

- Solid 5-nitro-2,3,4-trifluorobenzoic acid (0.75 g, 0.00339 mol) was dissolved in concentrated ammonium hydroxide (25 ml) to give instantly a yellow solution. A precipitate began to form within five minutes, after which time the mixture was acidified to pH 0 with concentrated aqueous hydrochloric acid. A yellow precipitate rapidly formed. The mixture was heated to boiling
- 30 and was filtered hot. The yellow solids were washed with 10 % aqueous hydrochloric acid and were suction dried to afford 0.47 g of a yellow powder; 64 % yield; <sup>1</sup>H-NMR (400 MHz; DMSO) δ 13.32 (s, 1H), 8.36 (d, 1H, J=7.6

Hz), 7.98 (s, 2H);  $^{19}\text{F}$ -NMR (376 MHz; DMSO)  $\delta$  -128.69 to -128.76 (m), -153.60 (d).

Step c: Preparation of methyl 4-amino-2,3-difluoro-5-nitrobenzoate

5 Hydrogen chloride gas was dissolved in anhydrous methanol (30 ml) until the solution was warm. The solid 4-amino-2,3-difluoro-5-nitrobenzoic acid (0.47 g; 0.00215 mol) was dissolved in this solution and the reaction mixture was brought to reflux with vigorous stirring for 23 hours under a nitrogen atmosphere. The reaction mixture was allowed to cool slowly on the  
10 bench. A yellow precipitate formed and was collected by vacuum filtration and dried with suction to afford 0.35 g of yellow microfilaments; 70 % yield; m.p. 183.5-184 °C;  $^1\text{H}$ -NMR (400 MHz; DMSO)  $\delta$  8.36 (dd, 1H,  $J=7.3, 1.7$  Hz), 8.06 (s, 2H), 3.78 (s, 3H);  $^{19}\text{F}$ -NMR (376 MHz; DMSO)  $\delta$  -128.85 to -128.92 (m), -153.29 (d); MS (APCI-) 231 (M-1, 100); IR (KBr) 3433, 3322, 1700,  
15 1650, 1549, 1343, 1285  $\text{cm}^{-1}$ ; Anal. calcd/found for:  $\text{C}_8\text{H}_6\text{F}_2\text{N}_2\text{O}_4$  C, 41.39/41.40; H, 2.61/2.50; N, 12.07/11.98; F, 16.37/16.58.

Step d: Preparation of methyl 4-amino-3-fluoro-2-(2-methyl-phenylamino)-5-nitrobenzoate

20 The solid methyl 4-amino-2,3-difluoro-5-nitrobenzoate (0.087 g,  $3.7 \times 10^{-4}$  mol) was dissolved in *ortho*-toluidine (3 ml, 0.028 mol). The reaction mixture was stirred at 200 °C for 35 minutes under a nitrogen atmosphere. The mixture was then partitioned between diethyl ether (150 ml) and 10 % aqueous hydrochloric acid (150 ml). The ether phase was dried with  
25 anhydrous magnesium sulfate and was concentrated *in vacuo* to a crude solid. The crude product was dissolved in 5 ml of dichloromethane and was filtered through a flash silica plug. Elution with dichloromethane afforded 0.0953 g of a yellow solid; 81 % yield; m.p. 164-168 °C;  $^1\text{H}$ -NMR (400 MHz; DMSO)  $\delta$  9.20 (s, 1H), 8.52 (d, 1H,  $J=1.7$  Hz), 7.57 (s, 2H), 7.19 (d, 1H,  $J=7.3$   
30 Hz), 7.12-7.08 (m, 1H), 7.02-6.98 (m, 1H), 6.95-6.91 (m, 1H), 3.78 (s, 3H), 2.21 (s, 3H);  $^{19}\text{F}$ -NMR (376 MHz; DMSO)  $\delta$  -141.13 (s); MS (APCI+) 320 (M+1, 100); (APCI-) 318 (M-1, 100); IR (KBr) 3467, 3346, 1690, 1305  $\text{cm}^{-1}$ ;



Anal. calcd/found for:  $C_{15}H_{14}FN_3O_4 \cdot 0.21 H_2O$  C, 55.77/55.97; H, 4.50/4.55; N, 13.01/12.61; F, 5.88/5.95.

Step e: Preparation of methyl 4,5-diamino-3-fluoro-2-(2-methyl-phenylamino)benzoate

To a mixture comprised of methyl 4-amino-3-fluoro-2-(2-methyl-phenylamino)-5-nitrobenzoate (2.52 g, 0.00789 mol), tetrahydrofuran (50 ml), methanol (50 ml) and washed Raney nickel (0.5 g) was initially applied 48.6 psi of hydrogen gas at 30.2 °C in a shaker for 4 hours 48 minutes. The mixture was filtered and the filtrate concentrated *in vacuo* to afford 2.20 g of a salmon-colored amorphous solid; 96 % yield;  $^1H$ -NMR (400 MHz; DMSO)  $\delta$  7.84 (s, 1H), 7.04 (d, 1H, J=7.1 Hz), 6.98 (d, 1H, J=1.2 Hz), 6.95-6.91 (m, 1H), 6.68-6.64 (m, 1H), 6.40-6.36 (m, 1H), 5.39 (s, 2H), 4.73 (s, 2H), 3.66 (s, 3H), 2.21 (s, 3H);  $^{19}F$ -NMR (376 MHz; DMSO)  $\delta$  -139.66 (s).

Step f: Preparation of methyl 7-fluoro-6-(2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylate

A stirring solution comprised of methyl 4,5-diamino-3-fluoro-2-(2-methyl-phenylamino)-benzoate (1.78 g, 0.00615 mol) in formic acid (Aldrich, 95-97 %, 100 ml, 2.5 mol) was brought to reflux for 3 hours followed by concentration *in vacuo* to give a crude brown solid. The crude product was triturated with chloroform (40 ml) and subsequently collected by vacuum filtration. The solids were dried with suction to afford 1.09 g of a light-lavender powder. The filtrate was concentrated *in vacuo* to a crude solid which was triturated with 10 ml of chloroform-dichloromethane. These solids were collected by vacuum filtration, rinsed with dichloromethane, and were suction-dried to give an additional 0.55 g of a light-lavender powder (total yield: 1.64 g); 87 % yield; m.p. 259-262 °C;  $^1H$ -NMR (400 MHz; DMSO)  $\delta$  8.42 (s, 1H), 8.03 (s, 1H), 7.93 (broad s, 1H), 7.12 (d, 1H, J=7.0 Hz), 6.99-6.95 (m, 1H), 6.75-6.71 (m, 1H), 6.48-6.44 (m, 1H), 3.81 (s, 3H), 2.30 (s, 3H);  $^{19}F$ -NMR (376 MHz; DMSO)  $\delta$  -132.84 (s); MS (APCI+) 300 (M+1, 100); (APCI-) 298 (M-1,

100); IR (KBr) 3322, 1689, 1437, 1326, 1218  $\text{cm}^{-1}$ ; Anal. calcd/found for:  $\text{C}_{16}\text{H}_{14}\text{FN}_3\text{O}_2 \cdot 0.32 \text{ H}_2\text{O}$  C, 62.99/63.01; H, 4.84/4.61; N, 13.77/13.70.

Step g: Preparation of methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylate

A stirring mixture comprised of methyl 7-fluoro-6-(2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylate (0.2492 g,  $8.326 \times 10^{-4}$  mol), benzyltrimethylammonium dichloriodinate (Aldrich, 95 %, 0.3934 g, 0.00113 mol), and zinc chloride (0.1899 g, 0.00139 mol) in glacial acetic acid (20 ml) was brought to reflux for 15 minutes. The hot suspension was filtered to isolate the precipitate which was dried in the vacuum oven (90  $^{\circ}\text{C}$ , ca. 10 mm Hg) overnight to afford 0.2392 g of a green powder; 68 % yield; m.p. 219-220  $^{\circ}\text{C}$  DEC;  $^1\text{H}$ -NMR (400 MHz; DMSO)  $\delta$  8.71 (s, 1H), 8.02 (s, 1H), 7.85 (broad s, 1H), 7.43 (d, 1H,  $J=1.7$  Hz), 7.24 (dd, 1H,  $J=8.5, 2.2$  Hz), 6.24 (dd, 1H,  $J=8.5, 5.4$  Hz), 3.76 (s, 3H), 2.22 (s, 3H);  $^{19}\text{F}$ -NMR (376 MHz; DMSO)  $\delta$  -132.86 (s); MS (APCI+) 426 (M+1, 48), 169 (100); (APCI-) 424 (M-1, 100); IR (KBr) 1704, 1508, 1227  $\text{cm}^{-1}$ .

Step h: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid

To a stirring solution comprised of methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylate (0.2035 g,  $4.786 \times 10^{-4}$  mol) in tetrahydrofuran (20 ml) was added solid potassium trimethylsilanolate (0.315 g, 0.00246 mol). The reaction mixture was stirred at ambient temperature under argon for 16 hours. An additional 0.082 g ( $6.39 \times 10^{-4}$  mol) of potassium trimethylsilanolate was added and the mixture stirred 30 minutes. The reaction mixture was concentrated *in vacuo* to one-third volume and was treated with diethyl ether (50 ml). The off-white precipitate formed was collected by vacuum filtration, giving a hygroscopic solid. The wet solid was dissolved in a 4:1 (v/v) ethyl acetate-methanol solution (500 ml). The solution was washed with 0.84 M aqueous citric acid (50 ml), dried ( $\text{MgSO}_4$ ), and concentrated *in vacuo* to a yellow liquid. The liquid was redissolved in

fresh ethyl acetate-methanol. The solution was washed with brine, dried (MgSO<sub>4</sub>), and concentrated *in vacuo*. The residue was redissolved in chloroform and reconcentrated to afford 1.55 g of a viscous yellow residue which was comprised mainly of citric acid; MS (APCI-) 191 (M-1, 100). The residue was dissolved in water (50 ml). Insoluble material was extracted into 1:1 (v/v) ethyl acetate-diethyl ether (250 ml). Upon separation, the aqueous phase remained strongly acidic (pH 0). The organic phase was washed with a fresh portion of water (150 ml). Upon separation, this wash was only slightly acidic (pH 4.5). The organic phase was dried (MgSO<sub>4</sub>), concentrated *in vacuo*, and chased with chloroform to give a tan semisolid. The product was triturated with hexanes. Vacuum filtration and suction-drying afforded 0.0839 g of a tan powder. A portion of the product (0.050 g) was recrystallized from boiling ethanol (1 ml). While cooling and moderate scratching, an off-white solid formed. This product was isolated by vacuum filtration and dried under high vacuum (23 °C) to afford 0.018 g of an off-white powder; 9 % yield; m.p. 247-248 °C DEC; <sup>19</sup>F-NMR (376 MHz; DMSO)  $\delta$  -132.87 (s); MS (APCI+) 412 (M+1, 100); (APCI-) 410 (M-1, 100); IR (KBr) 3322, 1689, 1437, 1326, 1218 cm<sup>-1</sup>; Anal. calcd/found for: C<sub>15</sub>H<sub>11</sub>FIN<sub>3</sub>O<sub>2</sub> · 0.61 C<sub>2</sub>H<sub>6</sub>O · 0.59 H<sub>2</sub>O (91.4 % parent) C, 43.30/43.30; H, 3.55/3.34; N, 9.34/9.15.

## EXAMPLE 2

Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide (PD 254552) (APK IC<sub>50</sub> < 10 nM (n = 2); colon 26 cells, 1 hour pretreatment, IC<sub>50</sub> = 20 nM)

Step a: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid pentafluorophenyl ester (PD 254551) (APK IC<sub>50</sub> = 120 nM (n=2))

To a stirring suspension comprised of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid (0.844 g, 2.05x10<sup>-3</sup> mol) in ethyl acetate (4 ml) was added a solution comprised of pentafluorophenol (0.375 g, 2.04x10<sup>-3</sup> mol) in N,N-dimethylformamide (10 ml). Solid

dicyclohexylcarbo-diimide (0.415 g,  $1.99 \times 10^{-3}$  mol) was then added and the reaction mixture was stirred for 22 hours. The reaction mixture was vacuum filtered to remove the precipitate that had formed. The filtrate was diluted with ethyl acetate (400 ml), and that solution was washed with water (3x400 ml),  
5 was dried ( $\text{MgSO}_4$ ), and was concentrated *in vacuo* to afford 1.7 g of a yellow foam. The crude product was purified by flash silica column chromatography. Elution with a gradient ( $\text{CHCl}_3$  to 0.5 % methanol in  $\text{CHCl}_3$ ) afforded 0.69 g of the yellow amorphous product; 60 % yield;  $^1\text{H-NMR}$  (400 MHz;  $\text{CDCl}_3$ )  $\delta$  8.54 (s, 1H), 8.28 (s, 1H), 8.04 (s, 1H), 7.49 (d, 1H,  $J=1.7$  Hz), 7.36 (dd, 1H,  $J=8.2$ ,  
10 1.7 Hz), 6.57 (dd, 1H,  $J=8.4$ , 6.5 Hz), 2.31 (s, 3H);  $^{19}\text{F-NMR}$  (376 MHz;  $\text{CDCl}_3$ )  $\delta$  -132.02 (s), -152.35 (d,  $J=18.3$  Hz), -157.26 (t,  $J=21.4$  Hz), -161.96 (dd,  $J=21.3$ , 18.3 Hz); MS (APCI+) 578 (M+1, 57), 394 (100); (APCI-) 576 (M-1, 44), 409 (100), 393 (95), 392 (82), 378 (55), 183 (97), 165 (68), 127 (53); IR (KBr)  $1731\text{ cm}^{-1}$  (C=O stretch).

15 Step b: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide

To a stirring solution comprised of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid pentafluorophenyl ester  
20 (0.63 g,  $1.09 \times 10^{-3}$  mol) in anhydrous tetrahydrofuran (5 ml) was added solid cyclopropylmethoxylamine hydrochloride (0.14 g,  $1.13 \times 10^{-3}$  mol) and diisopropylethylamine (0.6 ml,  $3.4 \times 10^{-3}$  mol). The reaction mixture was stirred for one week. The solvent was removed and the evaporate was treated with 10 % aqueous hydrochloric acid (200 ml) and was extracted with diethyl ether  
25 (200 ml). A biphasic suspension resulted, and the precipitate was isolated by vacuum filtration. The crude product was recrystallized from absolute ethanol to afford 0.18 g of a green-yellow powder; 35 % yield; mp  $168-172\text{ }^\circ\text{C}$ ;  $^1\text{H-NMR}$  (400 MHz; DMSO)  $\delta$  11.48 (s, 1H), 8.37 (s, 1H), 7.50 (broad s, 1H), 7.45 (s, 1H), 7.24 (s, 1H), 7.07 (d, 1H,  $J=8.4$  Hz), 6.03-5.97 (m, 1H), 3.38 (d, 2H,  
30  $J=6.5$  Hz), 2.04 (s, 3H), 0.85-0.75 (m, 1H), 0.30-0.22 (m, 2H), 0.00 (s, 2H);  $^{19}\text{F-NMR}$  (376 MHz; DMSO)  $\delta$  -133.23 (s); MS (APCI+) 481 (M+1, 77), 409 (100); (APCI-) 480 (M, 22), 407 (100); IR (KBr)  $1659$ ,  $1632$ ,  $1493\text{ cm}^{-1}$ ; Anal.

calcd/found for:  $C_{19}H_{18}FIN_4O_2 \cdot 0.50 HCl$  (96.3 % parent) C, 45.78/45.74; H, 3.74/3.84; N, 11.24/10.88.

### EXAMPLE 3

5

#### Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid hydroxyamide

10

#### Step a: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide

15

A solution comprised of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid, O-(tetrahydro-2H-pyran-2-yl)-hydroxylamine (1.25 equiv.), benzotriazole-1-yl-oxy-*tris*-pyrrolidino-phosphonium hexafluorophosphate (1.25 equiv.), and diisopropylethylamine (3 equiv.) in 1:1 v/v tetrahydrofuran-dichloromethane is stirred for 30 minutes. The reaction mixture is concentrated *in vacuo* and the residue is purified by flash chromatography; elution with dichloromethane affords the desired product. The product may be recrystallized with an appropriate solvent like methanol if further purification is necessary.

20

#### Step b: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid hydroxyamide

25

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide is dissolved in an appropriate hydrogen chloride-saturated solvent like methanol or ethanol. Once homogeneous, the solution is concentrated *in vacuo* to give the desired product. The product may be triturated with an appropriate solvent like chloroform or dichloromethane if further purification is necessary.

30

## EXAMPLE 4

## Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide

5

Step a: Preparation of O-cyclopropylmethylhydroxylamine hydrochlorideStep i: Preparation of 2-cyclopropylmethoxy-isoindole-1,3-dione

- 10 To a stirring solution/suspension comprised of N-hydroxyphthalimide (Aldrich, 57.15 g, 339.8 mmol), cyclopropanemethanol (Aldrich, 25.10 g, 341.1 mmol), and triphenylphosphine ("DEAD," Aldrich, 91.0 g, 344 mmol) in 1.00 L of tetrahydrofuran under a nitrogen atmosphere and cooled to 6 °C (internal mixture temperature) with an ice-water bath was added diethyl
- 15 azodicarboxylate (Aldrich, 56 ml, 356 mmol) dropwise over 20 minutes via addition funnel. The reaction mixture temperature was kept below 20 °C during the addition. Following addition of the DEAD, the cold bath was removed and the reaction mixture was stirred for 15 hours. The mixture was concentrated to a paste under reduced pressure. Chloroform (ca. 300 ml)
- 20 was added and the mixture swirled to loosen all solids. Vacuum filtration removed the insolubles. The filtrate was likewise filtered to remove white precipitate that formed and to give a clear filtrate. Concentration under reduced pressure afforded a clear oil. Flash filtration through silica gel (100 % chloroform) gave filtrates containing unseparated product. These filtrates
- 25 were combined and concentrated under reduced pressure to afford 127.4 g of a clear oil. The oil was dissolved in absolute ethanol (400 ml) and the solution was refrigerated for two hours. A white crystalline solid had precipitated and was subsequently collected by vacuum filtration. The product was dried in the vacuum oven (60 °C) to afford 42.66 g (58 %) of the desired material; m.p. 71-
- 30 77 °C; <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub> signal offset to δ 6.96) δ 7.54-7.43 (m, 4H), 3.74 (d, 2H, J=7.6 Hz), 1.02-0.95 (m, 1H), 0.34-0.30 (m, 1H), 0.04-0.00 (m, 1H).

Step ii: Preparation of O-cyclopropylmethylhydroxylamine hydrochloride

To a stirring solution comprised of 2-cyclopropylmethoxy-isoindole-1,3-dione (42.64 g, 196.3 mmol) in 150 ml of dichloromethane under ambient conditions was carefully added methylhydrazine (Aldrich, 10.7 ml, 197 mmol). A white precipitate began to form almost instantly. After 15 minutes of vigorous stirring, the suspension was vacuum filtered. The filtrate was likewise filtered to remove additional precipitate. The resulting clear filtrate was concentrated carefully (volatile product) under reduced pressure to afford a clear liquid/solid mixture. The white solids were removed when an ether (200 ml) solution of the product was made and vacuum filtered. The filtrate was acidified with gaseous hydrogen chloride, affording instantly a white precipitate. Collection of the solid by vacuum filtration and vacuum-oven drying (55 °C) afforded 18.7 g (77 %) of the white powder product; m.p. 165-168 °C; <sup>1</sup>H-NMR (400 MHz; DMSO) δ 10.77 (broad s, 2H), 3.57 (d, 2H, J=7.3 Hz), 0.84-0.74 (m, 1H), 0.31-0.25 (m, 2H), 0.04-0.00 (m, 1H); <sup>13</sup>C-NMR (100 MHz; DMSO) δ 75.39, 5.52, 0.00.

Step b: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide

A solution comprised of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid, O-cyclopropylmethylhydroxylamine hydrochloride (1.25 equiv.), benzotriazole-1-yl-oxy-*tris*-pyrrolidino-phosphonium hexafluorophosphate (1.25 equiv.), and diisopropylethylamine (3 equiv.) in 1:1 v/v tetrahydrofuran-dichloromethane is stirred for 30 minutes. The reaction mixture is concentrated *in vacuo* and the residue is taken up into diethyl ether. The ether phase is washed with dilute aqueous hydrochloric acid, saturated aqueous sodium bicarbonate, and brine, is dried (MgSO<sub>4</sub>), and is concentrated *in vacuo* to afford the desired product. The product may be recrystallized with an appropriate solvent like methanol or chloroform if further purification is necessary.

## EXAMPLE 5

Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylic acid

5

Step a: Preparation of 5-nitro-2,3,4-trifluorobenzoic acid

Same as for Example 1, Step a.

Step b: Preparation of 2,3-difluoro-4-hydroxy-5-nitrobenzoic acid

10 The solid 5-nitro-2,3,4-trifluorobenzoic acid (1.00 g, 0.00452 mol) was dissolved in 10 wt. % aqueous sodium hydroxide solution. The mixture was clear deep orange. After standing under ambient conditions for several minutes, the mixture was quenched with concentrated aqueous hydrochloric acid until strongly acidic (pH 0). A white solid precipitated which was isolated  
15 by vacuum filtration and dried with suction to afford 0.40 g of an off-white solid. This solid was recrystallized from chloroform (20 ml) to afford 0.22 g of an off-white crystalline powder; 22 % yield; MS (APCI-) 218 (M-1, 100).

Step c: Preparation of methyl 2,3-difluoro-4-hydroxy-5-nitrobenzoate

20 Anhydrous hydrogen chloride gas was dissolved in anhydrous methanol (50 ml) until the solution was warm. The microcrystalline solid 2,3-difluoro-4-hydroxy-5-nitrobenzoic acid 0.22 g, 0.00100 mol) was dissolved in the methanolic hydrogen chloride solution. The stirring reaction mixture was brought to reflux under nitrogen for 16 hours. The mixture was concentrated  
25 *in vacuo* to give a white solid. The product was dried under high vacuum to afford 0.213 g of a white powder; 91 % yield; m.p. 108-109.5 °C; <sup>1</sup>H-NMR (400 MHz; DMSO) δ 8.25 (dd, 1H, J=7.7, 2.2 Hz), 3.83 (s, 3H); (CDCl<sub>3</sub>) δ 10.83 (s, 1H), 8.66 (dd, 1H, J=7.0, 2.2 Hz), 3.98 (s, 3H); <sup>19</sup>F-NMR (376 MHz; DMSO) δ -127.85 (s), -154.32 (d, J=19.8 Hz); (CDCl<sub>3</sub>) δ -118.31 to -118.37  
30 (m), -152.38 (d, J=18.3 Hz); MS (APCI-) 232 (M-1, 100); IR (KBr) 3264, 1731, 1640, 1546, 1307, 1286, 1160 cm<sup>-1</sup>.



Step d: Preparation of 1-adamantyl 4-carboxymethyl-2,3-difluoro-6-nitrophenyl carbonate

To a solution comprised of 1-adamantyl fluoroformate (2.0 M) and pyridine (2.0 M) in tetrahydrofuran is added a stirred solution comprised of methyl 2,3-difluoro-4-hydroxy-5-nitrobenzoate (0.96 equiv., 0.384 M) in anhydrous tetrahydrofuran at ambient temperature. The reaction mixture is stirred for 6 hours and the solvent is removed *in vacuo*. The residue is dissolved in dichloromethane. The organic solution is washed with dilute aqueous hydrochloric acid, dilute aqueous sodium carbonate, and water, is dried (MgSO<sub>4</sub>), and is concentrated *in vacuo* to give the desired product.

Step e: Preparation of 1-adamantyl 4-carboxymethyl-2-fluoro-3-(2-methyl-phenylamino)-6-nitrophenyl carbonate

The compound 1-adamantyl 4-carboxymethyl-2,3-difluoro-6-nitrophenyl carbonate is dissolved in excess *ortho*-toluidine. The reaction mixture is stirred at 200 °C for 6 hours. The mixture is allowed to cool and is dissolved in diethyl ether. The organic phase is washed with dilute aqueous hydrochloric acid, saturated aqueous sodium bicarbonate, and brine, is dried (MgSO<sub>4</sub>), and is concentrated *in vacuo* to afford the desired product. The product is purified by flash chromatography as necessary.

Step f: Preparation of methyl 3-fluoro-4-hydroxy-2-(2-methyl-phenylamino)-5-nitrobenzoate

The compound 1-adamantyl 4-carboxymethyl-2-fluoro-3-(2-methyl-phenylamino)-6-nitrophenyl carbonate is dissolved in excess trifluoroacetic acid at ambient temperature. The mixture is stirred for 20 minutes. The TFA is removed under reduced pressure. The residue is subjected to vacuum pump to remove adamantan-1-ol to give the desired product.

Step g: Preparation of methyl 5-amino-3-fluoro-4-hydroxy-2-(2-methyl-phenylamino)-benzoate

The compound methyl 3-fluoro-4-hydroxy-2-(2-methyl-phenylamino)-5-nitrobenzoate is treated as in Step e, Example 1.

Step h: Preparation of methyl 7-fluoro-6-(2-methyl-phenylamino)-1H-benzooxazole-5-carboxylate

The compound 5-amino-3-fluoro-4-hydroxy-2-(2-methyl-phenylamino)-benzoate is treated as in Step f, Example 1. The product may be recrystallized with an appropriate solvent like chloroform or ethanol if further purification is necessary.

Step i: Preparation of methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylate

A stirring mixture comprised of methyl 7-fluoro-6-(2-methyl-phenylamino)-1H-benzooxazole-5-carboxylate (0.042 M), benzyltrimethylammonium dichloriodinate (Aldrich, 95 %, 0.057 M, 1.36 equiv.), and zinc chloride (0.070 M, 1.67 equiv.) in glacial acetic acid is brought to reflux for 15 minutes. The mixture is concentrated *in vacuo* and the residue taken up into diethyl ether. The ether solution is washed with dilute aqueous hydrochloric acid, water, and brine, is dried (MgSO<sub>4</sub>), and is concentrated *in vacuo* to obtain the desired product. The product may be purified by recrystallization with an appropriate solvent like ethanol.

Step j: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylic acid

To a stirring solution comprised of methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylate (0.024 M) in tetrahydrofuran is added solid potassium trimethylsilanolate (5.14 equiv.). The reaction mixture is stirred at ambient temperature under argon for 16 hours. An additional equivalent of potassium trimethylsilanolate is added and the mixture stirred 30 minutes. The reaction mixture is concentrated *in vacuo* to give a residue that

is then taken up into 1:1 (v/v) ethyl acetate-diethyl ether. The organic phase is washed with dilute aqueous hydrochloric acid, water, and brine, is dried (MgSO<sub>4</sub>), is concentrated *in vacuo*, and chased with chloroform to give a crude product. Recrystallization from an appropriate solvent like ethanol gives the purified desired product.

### EXAMPLE 6

#### Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylic acid hydroxyamide

Step a: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylic acid is treated as in Step a, Example 2.

Step b: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylic acid hydroxyamide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide is treated as in Step b, Example 2.

### EXAMPLE 7

#### Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylic acid cyclopropylmethoxy-amide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzooxazole-5-carboxylic acid is treated as in Step b, Example 3.

## EXAMPLE 8

Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylic acid

5

Step a: Preparation of 5-nitro-2,3,4-trifluorobenzoic acid

Same as for Example 1, Step a.

Step b: Preparation of 2,3-difluoro-4-hydroxy-5-nitrobenzoic acid

10

Same as for Example 4, Step b.

Step c: Preparation of methyl 2,3-difluoro-4-hydroxy-5-nitrobenzoate

Same as for Example 4, Step c.

15

Step d: Preparation of 4-dimethylthiocarbamoyloxy-2,3-difluoro-5-nitrobenzoic acid methyl ester

20

25

30

A solution of methyl 2,3-difluoro-4-hydroxy-5-nitrobenzoate in N,N-dimethylformamide is treated with one molar equivalent of cesium carbonate and warmed to 85 °C for 30 minutes. The stirring mixture is then treated dropwise rapidly with a solution comprised of a slight excess of N,N-dimethylthiocarbamoyl chloride in N,N-dimethylformamide. The reaction mixture is stirred at room temperature for one hour, or may be warmed over a steam bath for one hour. The mixture is then poured into water and extracted with ethyl acetate. The organic phase is washed with 5 % aqueous sodium hydroxide, water, and brine, and is then dried with a drying agent like magnesium sulfate or sodium sulfate. The solvent is then removed *in vacuo* to give a crude product. The compound is purified by ordinary methods such as chromatography or crystallization from an appropriate solvent.

Step e: Preparation of 4-Dimethylthiocarbamoyloxy-3-fluoro-5-nitro-2-o-tolylamino-benzoic acid methyl ester

The compound 4-dimethylthiocarbamoyloxy-2,3-difluoro-5-nitro-benzoic acid methyl ester is dissolved in excess o-toluidine. The stirring mixture is brought to 200 °C for one hour. The mixture is then poured into 5 % aqueous hydrochloric acid. The aqueous mixture is extracted with diethyl ether. The organic phase is washed with water and brine, is dried over magnesium sulfate, and is concentrated *in vacuo*. The crude product is purified by ordinary methods such as chromatography or crystallization from an appropriate solvent.

Step f: Preparation of methyl 7-fluoro-6-(2-methyl-phenylamino)-1H-benzothiazole-5-carboxylate

The compound methyl 5-amino-3-fluoro-4-mercapto-2-(2-methyl-phenylamino)-benzoate is treated as in Step h, Example 4.

Step g: Preparation of methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylate

The compound methyl 7-fluoro-6-(2-methyl-phenylamino)-1H-benzothiazole-5-carboxylate is treated as in Step i, Example 4.

Step h: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylic acid

The compound methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylate is treated as in Step j, Example 4.

### EXAMPLE 9

Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylic acid hydroxyamide

Step a: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylic acid is treated as in Step a, Example 2.

Step b: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylic acid hydroxyamide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide is treated as in Step b, Example 2.

### EXAMPLE 10

Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylic acid cyclopropylmethoxy-amide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzothiazole-5-carboxylic acid is treated as in Step b, Example 3.

### EXAMPLE 11

Preparation of 8-fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid

Step a: Preparation of 8-fluoro-7-(2-methyl-phenylamino)-quinoxaline-6-carboxylic acid

The compound methyl 4,5-diamino-3-fluoro-2-(2-methyl-phenylamino)-benzoate (from Step e, Example 1) is dissolved in 2:1:1.2 v/v/v of 2.0 M acetic acid-4.0 M sodium acetate-methanol. The suspension is warmed to 65 °C (or until homogeneous) and the clear solution is poured into a 0.078 M aqueous sodium glyoxal bisulfite (Aldrich, monohydrate, 1.05 equiv.) solution which is warmed to 70 °C. The reaction mixture is stirred gently between 55-75 °C for one hour, and is then cooled to 12 °C with an ice-water bath. Pulverized

sodium hydroxide pellets (27 equiv.) are added to the cold solution. The mixture is gently warmed to 30 °C and stirred for 45 minutes. The temperature is raised to 70 °C for 15 minutes. The mixture is allowed to cool and is treated with ethyl acetate. The biphasic mixture is treated with concentrated aqueous hydrochloric acid to achieve pH 0 in the aqueous phase. The organic phase is separated, dried (MgSO<sub>4</sub>), and concentrated *in vacuo* to give the desired product. The product may be triturated with an appropriate solvent like dichloromethane or recrystallized from a solvent like ethanol for further purification as necessary.

Step b: Preparation of 8-fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid

The compound 8-fluoro-7-(2-methyl-phenylamino)-quinoxaline-6-carboxylic acid is treated as in Step i, Example 4.

EXAMPLE 12

Preparation of 8-fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid hydroxyamide

Step a: Preparation of 8-fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide

The compound 8-fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid is treated as in Step a, Example 2.

Step b: Preparation of 8-fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid hydroxyamide

The compound 8-fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide is treated as in Step b, Example 2.

## EXAMPLE 13

Preparation of 8-fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid cyclopropylmethoxy-amide

- 5        The compound 8-fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid is treated as in Step b, Example 3.

## EXAMPLE 14

10    Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic acid

Step a: Preparation of methyl 7-fluoro-6-(2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylate

- 15    To a stirring solution comprised of methyl 4,5-diamino-3-fluoro-2-(2-methyl-phenylamino)-benzoate (from Step e, Example 1) and diisopropylethylamine (2 equiv.) in an appropriate solvent like diethyl ether or toluene is added a reagent like N-thioaniline or thionyl chloride (1.35 equiv.). The reaction mixture is brought to reflux for one hour. The mixture is quenched with dilute aqueous hydrochloric acid. The organic phase is washed with saturated aqueous sodium bicarbonate and brine, is dried (MgSO<sub>4</sub>), and is concentrated *in vacuo* to afford the desired product. The product may be recrystallized with an appropriate solvent like chloroform or ethanol, or may be chromatographed if further purification is necessary.

25

Alternative method: The compound methyl 4,5-diamino-3-fluoro-2-(2-methyl-phenylamino)-benzoate is added to a stirring solution of sulfur monochloride (6 equiv.) in N,N-dimethylformamide and the mixture is gradually heated to 75-80 °C. After 5 hours the mixture is cooled to 10 °C, water is slowly added.

30

The mixture is extracted with a solvent like diethyl ether or dichloromethane. The organic extract is dried (MgSO<sub>4</sub>) and is concentrated *in vacuo* to afford the desired product. The product may be recrystallized with an appropriate



solvent like chloroform or ethanol, or may be chromatographed if further purification is necessary.

Step b: Preparation of methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylate

The compound methyl 7-fluoro-6-(2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylate is treated as in Step i, Example 4.

Step c: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic acid

The compound methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylate is treated as in Step j, Example 4.

#### EXAMPLE 15

Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic acid hydroxyamide

Step a: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic acid is treated as in Step a, Example 2.

Step b: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic acid hydroxyamide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide is treated as in Step b, Example 2.

## EXAMPLE 16

Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-  
benzo[1,2,5]thiadiazole-5-carboxylic acid cyclopropylmethoxy-amide

5        The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-  
benzo[1,2,5]thiadiazole-5-carboxylic acid is treated as in Step b, Example 3.

## EXAMPLE 17

10    Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-  
benzo[1,2,5]oxadiazole-5-carboxylic acid

Step a: Preparation of methyl 7-fluoro-6-(2-methyl-phenylamino)-  
benzo[1,2,5]oxadiazole-5-carboxylate 2-oxide

15        See Takakis, I. M.; Hadjimihalakis, P. M., *J. Heterocyclic Chem.*, **27**,  
177 (1990).

A mixture comprised of methyl 4-amino-3-fluoro-2-(2-methyl-  
phenylamino)-5-nitrobenzoate (from Step d, Example 1) and  
iodosobenzenediacetate (1.76 equiv.) in benzene is stirred at ambient  
20    temperature for 5 hours. The mixture is concentrated *in vacuo* and the  
residue purified by column chromatography to give the desired product.

Alternative method: A solution comprised of methyl 4-amino-3-fluoro-2-(2-  
methyl-phenylamino)-5-nitrobenzoate (0.86 M) in tetrahydrofuran is diazotized  
25    and the diazonium salt is treated *in situ* with sodium azide as described by  
Smith, P. A. S.; Boyer, J. H., *Org. Synth.*, **31**, 14 (1951) and references 4 and  
8 cited therein. Thermolysis of this intermediate in ethylene glycol at 110-120  
°C for one hour affords the desired product.

30    Step b: Preparation of methyl 7-fluoro-6-(2-methyl-phenylamino)-  
benzo[1,2,5]oxadiazole-5-carboxylate

A solution comprised of methyl 7-fluoro-6-(2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylate 2-oxide and sodium azide (1.38 equiv.) in ethylene glycol is heated to 140-150 °C for 30 minutes to obtain, after column chromatography, the desired product.

Step c: Preparation of methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylate

The compound methyl 7-fluoro-6-(2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylate is treated as in Step i, Example 4.

Step d: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylic acid

The compound methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylate is treated as in Step j, Example 4.

### EXAMPLE 18

Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylic acid hydroxyamide

Step a: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylic acid is treated as in Step a, Example 2.

Step b: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylic acid hydroxyamide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide is treated as in Step b, Example 2.

## EXAMPLE 19

Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-  
benzo[1,2,5]oxadiazole-5-carboxylic acid cyclopropylmethoxy-amide

5        The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-  
benzo[1,2,5]oxadiazole-5-carboxylic acid is treated as in Step b, Example 3.

## EXAMPLE 20

10    Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-  
carboxylic acid

Step a: Preparation of methyl 7-fluoro-6-(2-methyl-phenylamino)-1H-  
benzotriazole-5-carboxylate

15        The compound methyl 4,5-diamino-3-fluoro-2-(2-methyl-phenylamino)-  
benzoate (from Step e, Example 1) is diazotized by ordinary methods.  
Workup gives the desired product.

20    Step b: Preparation of methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-  
benzotriazole-5-carboxylate

The compound methyl 7-fluoro-6-(2-methyl-phenylamino)-1H-  
benzotriazole-5-carboxylate is treated as in Step i, Example 4.

25    Step c: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-  
benzotriazole-5-carboxylic acid

The compound methyl 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-  
benzotriazole-5-carboxylate is treated as in Step j, Example 4.

## EXAMPLE 21

30    Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-  
carboxylic acid hydroxyamide

Step a: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-carboxylic acid is treated as in Step a, Example 2.

Step b: Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-carboxylic acid hydroxyamide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-carboxylic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide is treated as in Step b, Example 2.

## EXAMPLE 22

Preparation of 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-carboxylic acid cyclopropylmethoxy-amide

The compound 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-carboxylic acid is treated as in Step b, Example 3.


## F. OTHER EMBODIMENTS

From the above disclosure and examples, and from the claims below, the essential features of the invention are readily apparent. The scope of the invention also encompasses various modifications and adaptations within the knowledge of a person of ordinary skill. Examples include a disclosed compound modified by addition or removal of a protecting group, or an ester, pharmaceutical salt, hydrate, acid, or amide of a disclosed compound. Publications cited herein are hereby incorporated by reference in their entirety.

What is claimed is:

CLAIMS

1. A method for treating chronic pain, said method comprising administering to a subject in need of such treatment a composition comprising a MEK inhibitor selected from a compound of the following formula (I):



(I)

wherein

W is  $OR_1$ ,  $NR_2OR_1$ ,  $NR_AR_B$ ,  $NR_2NR_AR_B$ ,  $O(CH_2)_{2-4}NR_AR_B$ , or  $NR_2(CH_2)_{2-4}NR_AR_B$ ;

$R_1$  is H, C<sub>1-8</sub> alkyl, C<sub>3-8</sub> alkenyl, C<sub>3-8</sub> alkynyl, C<sub>3-8</sub> cycloalkyl, phenyl, (phenyl)-C<sub>1-4</sub> alkyl, (phenyl)C<sub>3-4</sub> alkenyl, (phenyl)C<sub>3-4</sub> alkynyl, (C<sub>3-8</sub> cycloalkyl)C<sub>1-4</sub> alkyl, (C<sub>3-8</sub> cycloalkyl)C<sub>3-4</sub> alkenyl, (C<sub>3-8</sub> cycloalkyl)C<sub>3-4</sub> alkynyl, C<sub>3-8</sub> heterocyclic radical, (C<sub>3-8</sub> heterocyclic radical)C<sub>1-4</sub> alkyl, (C<sub>3-8</sub> heterocyclic radical)-C<sub>3-4</sub> alkenyl, (C<sub>3-8</sub> heterocyclic radical)C<sub>3-4</sub> alkynyl or  $(CH_2)_{2-4}NR_CR_D$ ;

$R_2$  is H, C<sub>1-4</sub> alkyl, phenyl, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> heterocyclic radical, or (C<sub>3-6</sub> cycloalkyl) methyl;


$R_A$  is H, C<sub>1-6</sub> alkyl, C<sub>3-8</sub> alkenyl, C<sub>3-8</sub> alkynyl, C<sub>3-8</sub> cycloalkyl, phenyl, (C<sub>3-8</sub> cycloalkyl)C<sub>1-4</sub> alkyl, (C<sub>3-8</sub> cycloalkyl)C<sub>3-4</sub> alkenyl, (C<sub>3-8</sub> cycloalkyl)C<sub>3-4</sub> alkynyl, C<sub>3-8</sub> heterocyclic radical, (C<sub>3-8</sub> heterocyclic radical)C<sub>1-4</sub> alkyl, (aminosulfonyl)phenyl, [(aminosulfonyl)phenyl]C<sub>1-4</sub> alkyl, (aminosulfonyl)C<sub>1-6</sub>

alkyl, (aminosulfonyl)C<sub>3-6</sub> cycloalkyl, [(aminosulfonyl)C<sub>3-6</sub> cycloalkyl]C<sub>1-4</sub> alkyl, or (CH<sub>2</sub>)<sub>2-4</sub> NR<sub>C</sub>R<sub>D</sub>;


R<sub>B</sub> is H, C<sub>1-8</sub> alkyl, C<sub>3-8</sub> alkenyl, C<sub>3-8</sub> alkynyl, C<sub>3-8</sub> cycloalkyl, or phenyl;

5


Q is one of the following formulae (i) – (iii):



(i)



(ii)



(iii)


10 R<sub>3</sub> is H or F;

R<sub>4</sub> is halo, NO<sub>2</sub>, SO<sub>2</sub>NR<sub>O</sub>(CH<sub>2</sub>)<sub>2-4</sub>NR<sub>E</sub>R<sub>F</sub>, SO<sub>2</sub>NR<sub>E</sub>R<sub>F</sub>, or (CO)T;


T is C<sub>1-8</sub> alkyl, C<sub>3-8</sub> cycloalkyl, (NR<sub>E</sub>R<sub>F</sub>)C<sub>1-4</sub> alkyl, OR<sub>F</sub>, -NR<sub>O</sub>(CH<sub>2</sub>)<sub>2-4</sub>NR<sub>E</sub>R<sub>F</sub>,

15 or NR<sub>E</sub>R<sub>F</sub>;


Z is one of the following formulae (iv) – (viii):



(iv)




(v)




(vi)

20



(vii)



(viii)

one of  $R_5$  and  $R_6$  is H or methyl and the other of  $R_5$  and  $R_6$  is H,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl, phenyl, benzyl, or  $-M-E-G$ ;

M is O, CO,  $SO_2$ ,  $NR_J$ ,  $(CO)NR_H$ ,  $NR_H(CO)$ ,  $NR_H(SO_2)$ ,  $(SO_2)NR_H$ , or  $CH_2$ ;

E is  $(CH_2)_{1-4}$  or  $(CH_2)_m O(CH_2)_p$  where  $1 \leq (\text{each of } m \text{ and } p) \leq 3$  and  $2 \leq (m + p) \leq 4$ ; or E is absent;

G is  $R_K$ ,  $OR_I$  or  $NR_JR_K$ , provided that if  $p = 1$ , then G is H;

$R_7$  is H,  $C_{1-4}$  alkyl,  $C_{2-4}$  alkenyl,  $C_{2-4}$  alkynyl,  $C_{3-6}$  cycloalkyl, phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl,  $(CH_2)_{1-2}Ar$ , where Ar is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl,

$SO_2NR_H(CH_2)_{2-4}NR_JR_K$ ,  $(CO)(CH_2)_{2-4}NR_JR_K$  or  $(CO)NR_H(CH_2)_{2-4}NR_JR_K$ ;

$X_1$  is O, S,  $NR_8$ , or  $CHR_9$ ;  $X_2$  is O, S, or  $CHR_9$ ; and  $X_3$  is O or S; where if  $X_1$  or  $X_2$  is  $CHR_9$ , said compound may also be a tautomerized indole;

$R_8$  is H,  $C_{1-4}$  alkyl, phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl,  $(CH_2)_{1-2}Ar$ , where Ar is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl,  $C_{2-4}$  alkenyl,  $C_{2-4}$  alkynyl,  $C_{3-6}$  cycloalkyl, or  $(C_{2-4} \text{ alkyl})NR_LR_M$ ; provided  $R_7$  and  $R_8$  together have no more than 14 carbon atoms, exclusive of  $R_L$ ,  $R_M$ ,  $R_J$  and  $R_K$ ;



$R_G$  is  $C_{1-4}$  alkyl, phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl,  $C_{3-4}$  alkenyl,  $C_{3-4}$  alkynyl,  $C_{3-6}$  cycloalkyl,  $(CO)OR_P$ ,  $(C_{2-4}$  alkyl) $NR_LR_M$ ,  $(CO)NR_N(CH_2)_{2-4}NR_LR_M$ ,  $(CO)NR_LR_M$ ,  $(CO)(CH_2)_{2-4}-NR_LR_M$ , or  $(CH_2)_{1-2}Ar$ , where  $Ar$  is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl;

5

$R_9$  is  $C_{1-4}$  alkyl, phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl,  $C_{2-4}$  alkenyl,  $C_{2-4}$  alkynyl,

$C_{3-6}$  cycloalkyl,  $(CO)OR_P$ ,  $(C_{2-4}$  alkyl) $NR_LR_M$ ,  $(CO)NR_N(CH_2)_{2-4}NR_LR_M$ ,  $(CO)NR_LR_M$ ,  $(CO)(CH_2)_{2-4}-NR_LR_M$ , or  $(CH_2)_{1-2}Ar'$ , where  $Ar'$  is phenyl, 2-

10

pyridyl, 3-pyridyl, or 4-pyridyl;

$R_P$  is H,  $C_{1-6}$  alkyl, phenyl,  $C_{3-4}$  alkenyl,  $C_{3-4}$  alkynyl,  $C_{3-6}$  cycloalkyl, or  $(CH_2)_{2-4}NR_LR_M$ ;

15

$R_{10}$  is H, methyl, halo, or  $NO_2$ ;

$R_{11}$  is H, methyl, halo, or  $NO_2$ ;

each of  $R_C$ ,  $R_D$ ,  $R_E$ ,  $R_F$ ,  $R_I$ ,  $R_J$ ,  $R_K$ ,  $R_L$  and  $R_M$  is independently selected from H,

20

$C_{1-4}$  alkyl,  $C_{3-4}$  alkenyl,  $C_{3-4}$  alkynyl,  $C_{3-6}$  cycloalkyl, and phenyl; each of  $NR_CR_D$ ,  $NR_ER_F$ ,  $NR_JR_K$ , and  $NR_LR_M$  can also independently be morpholinyl, piperazinyl, pyrrolidinyl, or piperadinyl; and

25

each of  $R_H$ ,  $R_N$ , and  $R_O$  is independently H, methyl, or ethyl;

wherein each hydrocarbon radical or heterocyclic radical above is optionally substituted with between 1 and 3 substituents independently selected from halo,  $C_{1-4}$  alkyl,  $C_{3-6}$  cycloalkyl,  $C_{3-4}$  alkenyl,  $C_{3-4}$  alkynyl, phenyl, hydroxyl, amino, (amino)sulfonyl, and  $NO_2$ , wherein each substituent alkyl, cycloalkyl, alkenyl, alkynyl or phenyl is in turn optionally substituted with between 1 and 3

30

substituents independently selected from halo, C<sub>1-2</sub> alkyl, hydroxyl, amino, and NO<sub>2</sub>;

or a pharmaceutically acceptable salt or C<sub>1-7</sub> ester thereof.

5

2. The method of claim 1, wherein said chronic pain is selected from neuropathic pain, idiopathic pain, and pain associated with chronic alcoholism, vitamin deficiency, uremia, or hypothyroidism.

10

3. The method of claim 2, wherein said chronic pain is a type of neuropathic pain.

15

4. The method of claim 3, wherein said neuropathic pain is associated with one of the following: inflammation, postoperative pain, phantom limb pain, burn pain, gout, trigeminal neuralgia, acute herpetic and postherpetic pain, causalgia, diabetic neuropathy, plexus avulsion, neuroma, vasculitis, viral infection, crush injury, constriction injury, tissue injury, limb amputation, post-operative pain, arthritis pain, and any other nerve injury between the peripheral nervous system and the central nervous system, inclusively.

20

5. The method of claim 2, wherein said chronic pain is associated with chronic alcoholism, vitamin deficiency, uremia, or hypothyroidism.

25

6. The method of claim 2, wherein said chronic pain is associated with idiopathic pain.

7. The method of claim 1, wherein said chronic pain is associated with inflammation.

30

8. The method of claim 1, wherein said chronic pain is associated with arthritis.

9. The method of claim 1, wherein said chronic pain is associated with post-operative pain.

5 10. A method of claim 1, wherein Q is formula (i).

11. A method of claim 10, wherein  $R_3$  is H or fluoro.

12. A method of claim 11, wherein  $R_4$  is fluoro, chloro, or bromo.

10

13. A method of claim 1, wherein  $R_{10}$  is hydrogen, methyl, fluoro, or chloro.

15

14. A method of claim 1, wherein  $R_{11}$  is methyl, chloro, fluoro, nitro, or hydrogen.

15. A method of claim 14, wherein  $R_{11}$  is H.

16. A method of claim 14, wherein  $R_{11}$  is fluoro.

20

17. A method of claim 13, wherein each of  $R_{10}$  and  $R_{11}$  is fluoro.

25

18. A method of claim 1, wherein  $R_1$  is H, methyl, ethyl, propyl, isopropyl, isobutyl, benzyl, phenethyl, allyl,  $C_{3-5}$  alkenyl,  $C_{3-6}$  cycloalkyl,  $(C_{3-5}$  cycloalkyl) $C_{1-2}$  alkyl,  $(C_{3-5}$  heterocyclic radical) $C_{1-2}$  alkyl, or  $(CH_2)_{2-4} NR_C R_D$ .

19. A method of claim 18, wherein  $R_1$  is H or  $(C_{3-4}$  cycloalkyl) $C_{1-2}$  alkyl.

30

20. A method of claim 1, wherein  $R_2$  is H or methyl.

21. A method of claim 1, wherein  $R_A$  has at least one hydroxyl substituent.

22. A compound of claim 1, wherein  $R_A$  is H, methyl, ethyl, isobutyl, hydroxyethyl, phenyl, 2-piperidin-1-yl-ethyl, 2,3-dihydroxy-propyl, 3-[4-(2-hydroxyethyl)-piperazin-1-yl]-propyl, 2-pyrrolidin-1-yl-ethyl, or 2-diethylamino-ethyl; and  $R_B$  is H; or where  $R_B$  is methyl and  $R_A$  is phenyl.

23. A method of claim 1, wherein W is  $NR_AR_B$  or  $NR_2NR_AR_B$ .

24. A method of claim 1, wherein W is  $NR_2(CH_2)_{2-4}NR_AR_B$  or  $O(CH_2)_{2-3}NR_AR_B$ .

25. A method of claim 1, wherein W is  $NR_2OR_1$ .

26. A method of claim 1, wherein W is  $OR_1$ .

27. A method of claim 1, wherein Z is formula (v).

28. A method of claim 27, wherein  $X_1$  is  $NR_8$ , and  $R_7$  is H.

29. A method of claim 1, wherein said MEK inhibitor has a structure selected from: 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzimidazole-5-carboxylic acid.

30. A method of claim 1, wherein said MEK inhibitor has a structure selected from: 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzimidazole-5-carboxylic acid; 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzooxazole-5-carboxylic acid; 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzothiazole-5-carboxylic acid; 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]thiadiazole-5-carboxylic acid; 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-benzo[1,2,5]oxadiazole-5-carboxylic acid; 7-fluoro-6-(4-iodo-2-

5 methyl-phenylamino)-2-(2-hydroxyethyl)-1H-benzoimidazole-5-carboxylic acid; 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-2-(2-dimethylamino-ethyl)-1H-benzoimidazole-5-carboxylic acid; 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1-acetyl-benzoimidazole-5-carboxylic acid; 8-fluoro-7-(4-iodo-2-methyl-phenylamino)-quinoxaline-6-carboxylic acid; and 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzotriazole-5-carboxylic acid; and the corresponding hydroxamic acids and cyclopropylmethyl hydroxamates.

10 31. The method of claim 1 wherein said MEK inhibitor has a structure selected from: 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide; 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-6,7-dihydro-1H-benzoimidazole-5-carboxylic acid (hydrochloride); 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid; 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-15 3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide; 6-(2-chloro-4-iodo-phenylamino)-7-fluoro-1H-benzoimidazole-5-carboxylic acid; and 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid pentafluorophenyl ester.

20 32. The method of claim 1 wherein said MEK inhibitor has a structure selected from: 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-1H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide; and 7-fluoro-6-(4-iodo-2-methyl-phenylamino)-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide.

25

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 January 2001 (25.01.2001)

PCT

(10) International Publication Number  
**WO 01/05390 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 31/404**,  
31/4192, 31/4245, 31/433, 31/42, 31/428, 31/498, A61P  
25/04

Robert, Denham [GB/GB]; 3 Teasel Way, Cambridge  
CB1 9YT (GB).

(21) International Application Number: PCT/US00/18345

(74) Agents: RYAN, M., Andrea; Warner-Lambert Company,  
201 Tabor Road, Morris Plains, NJ 07950 et al. (US).

(22) International Filing Date: 5 July 2000 (05.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(81) Designated States (*national*): AE, AG, AL, AU, BA, BB,  
BG, BR, BZ, CA, CN, CR, CU, CZ, DM, DZ, EE, GD, GE,  
HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV,  
MA, MG, MK, MN, MX, MZ, NO, NZ, PL, RO, SG, SI,  
SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA.

(30) Priority Data:  
60/144,418 16 July 1999 (16.07.1999) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):  
**WARNER-LAMBERT COMPANY** [US/US]; 201  
Tabor Road, Morris Plains, NJ 07950 (US).

**Published:**

— With international search report.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BARRETT**,  
Stephen, Douglas [US/US]; 14220 Sunbury, Livonia, MI  
48154 (US). **BRIDGES**, Alexander, James [GB/US];  
3301 Textile Road, Saline, MI 48176 (US). **TECLE**, Haile  
[US/US]; 3048 Turnberry, Ann Arbor, MI 48108 (US).  
**DIXON**, Alistair [GB/GB]; 108 Gwydir Street, Cam-  
bridge CB1 2LL (GB). **LEE**, Kevin [GB/GB]; 81 Williams  
Smith Close, Cambridge CB1 9YT (GB). **PINNOCK**,

(88) Date of publication of the international search report:  
17 May 2001

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

WO 01/05390 A3


(54) Title: METHOD FOR TREATING CHRONIC PAIN USING MEK INHIBITORS

(57) Abstract: The invention features a method for treating chronic pain using a compound of formula (I) shown in claim 1 of the application.

10/03/037

1/8

**FIG. 1** EFFECT OF PD 198306 ON STREPTOZOCIN-INDUCED STATIC ALLODYNIA




\*\* P<0.01 VS VEH TEST SAME DAY (Mann-Whitney) (n=6-7)

10/031037

2/8


**FIG. 2** EFFECT OF SYSTEMIC (p.o.) ADMINISTRATION OF PD 198306 ON STREPTOZOCIN-INDUCED STATIC ALLODYNIA





3/8


**FIG. 3** EFFECT OF SYSTEMIC (p.o.) ADMINISTRATION OF PD 198306 ON CCI-INDUCED STATIC ALLODYNIA



10/03/039

4/8


**FIG. 4** EFFECT OF INTRATHECAL (i.t.) ADMINISTRATION OF PD 198306 ON STREPTOZOCIN-INDUCED STATIC ALLODYNIA



19/03/037

5/8

**FIG. 5** EFFECT OF INTRATHECAL (i.t.) ADMINISTRATION OF PD 198306 ON CCI-INDUCED STATIC ALLODYNIA




10/03/037

6/8

**FIG. 6** EFFECT OF INTRAPLANTAR (i.pl.) ADMINISTRATION OF PD 198306 ON STREPTOZOCIN-INDUCED STATIC ALLODYNIA

□ VEHICLE (i.pl.)  
PD 198306  
■ i.pl. (3mg/100 $\mu$ l)  
▨ i.t. (30 $\mu$ g/10 $\mu$ l)




10/631037

7/8

**FIG. 7** EFFECT OF INTRAPLANTAR (i.pl.) ADMINISTRATION OF PD 198306 ON CCI-INDUCED STATIC ALLODYNIA


VEHICLE (i.pl.)  
PD 198306  
i.pl. (3mg/100 $\mu$ l)  
i.t. (30ug/10 $\mu$ l)



10/03/037

8/8

**FIG. 8** EFFECT OF INTRATHECAL (i.t.) ADMINISTRATION OF PD 219622, PD 297447, PD 184352, PD 254552 OR PREGABALIN ON CCI-INDUCED STATIC ALLODYNIA



Docket No.  
A0000102-01-SMH

## Declaration and Power of Attorney For Patent Application

### English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

METHOD FOR TREATING CHRONIC PAIN USING MEK INHIBITORS

the specification of which

(check one)

☐ is attached hereto.

☒ was filed on July 5, 2000 As United States Application No. \_\_\_\_\_ or PCT International  
Application Number PCT/US00/18345  
and was amended on \_\_\_\_\_

(if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Applications

Priority Not Claimed

(Number) (Country) (Day/Month/Year Filed)

☐

(Number) (Country) (Day/Month/Year Filed)

☐

(Number) (Country) (Day/Month/Year Filed)

☐

I hereby claim the benefit under 35 U.S.C. Section 119(e) of any United States provisional application(s) listed below:

60/144418	16 July 1999
(Application Serial No.)	(Filing Date)

(Application Serial No.)	(Filing Date)
--------------------------	---------------

(Application Serial No.)	(Filing Date)
--------------------------	---------------

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

PCT/US00/18345	05 July 2000	Pending
(Application Serial No.)	(Filing Date)	(Status)
		(patented, pending, abandoned)

(Application Serial No.)	(Filing Date)	(Status)
		(patented, pending, abandoned)

(Application Serial No.)	(Filing Date)	(Status)
		(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.



POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (*list name and registration number*)

Charles W. Ashbrook 27,610  
 Heidi M. Berven 48,951  
 Evan J. Federman 37,060  
 Mehdi Ganjeizadeh 47,585  
 Rosanne Goodman 32,534  
 Suzanne M. Harvey 42,640  
 David R. Kurlandsky 41,505

Darryl C. Little 40,703  
 J. Trevor Lumb 28,567  
 Claude F. Purchase, Jr. 47,871  
 James Proscia 47,010  
 Francis J. Tinney 33,069  
 Linda Vag 32,071

Send Correspondence to: Suzanne M. Harvey  
Warner-Lambert Company  
2800 Plymouth Road  
Ann Arbor, Michigan 48105

Direct Telephone Calls to: (*name and telephone number*)  
Suzanne M. Harvey, (734) 622-2658

Full name of sole or first inventor

STEPHEN DOUGLAS BARRETT

Sole or first inventor signature

Stephen Douglas Barrett

Date

11/13/01

Residence

Livonia, Michigan 48154 United States

Citizenship

United States

Post Office Address

14220 Sunbury

Livonia, Michigan 48154 United States

Full name of sole or second inventor

ALEXANDER JAMES BRIDGES

Sole or second inventor signature

Alexander James Bridges

Date

11/2/01

Residence

Saline, Michigan 48176 United States

Citizenship

Great Britain

Post Office Address

3301 Textile Road

Saline, Michigan 48176 United States

Full name of sole or third inventor

HAILE TECLE

Sole or third inventor signature

*Haile Tecle*

Date

11/07/01

Residence

Ann Arbor, Michigan 48108 United States

Citizenship

United States *MT*

Post Office Address

3048 Turnberry

Ann Arbor, Michigan 48108 United States

Full name of sole or fourth inventor

ALISTAIR DIXON

Sole or fourth inventor signature

Date

SEE ATTACHED

Residence

Cambridge, UK CB1 2LL *CB1*

Citizenship

Great Britain

Post Office Address

108 Gwydir Street

Cambridge, UK CB1 2LL

Full name of sole or fifth inventor

KEVIN LEE

Sole or fifth inventor signature

*Kevin Lee*

Date

23/12/01

Residence

Cambridge, UK CB1 9YT *Kenilworth, UK CV8 1D5*

Citizenship

Great Britain

Post Office Address

81 William Smith Close *ll**S Archer Road*

Cambridge, UK CB1 9YT

*Kenilworth, UK CV8 1D5*

Full name of sole or sixth inventor

ROBERT PINNOCK

Sole or sixth inventor signature

*Robert Pinnock*

Date

13th Dec 01

Residence

Cambridge, UK CB1 9YT *CB1*

Citizenship


Great Britain

Post Office Address

3 Teasel Way

Cambridge, UK CB1 9YT

Full name of sole or third inventor	
HAILE TECLE	
Sole or third inventor signature	Date
Residence	
Ann Arbor, Michigan 48108 United States	
Citizenship	
United States	
Post Office Address	
3048 Turnberry	
Ann Arbor, Michigan 48108 United States	

Full name of sole or fourth inventor	
ALISTAIR DIXON	
Sole or fourth inventor signature	Date
	8/5/02
Residence	
Cambridge, UK CB1 2LL	
Citizenship	
Great Britain	
Post Office Address	
108 Gwydir Street	
Cambridge, UK CB1 2LL	

Full name of sole or fifth inventor	
KEVIN LEE	
Sole or fifth inventor signature	Date
Residence	
Cambridge, UK CB1 9YT	
Citizenship	
Great Britain	
Post Office Address	
81 William Smith Close	
Cambridge, UK CB1 9YT	

Full name of sole or sixth inventor	
ROBERT PINNOCK	
Sole or sixth inventor signature	Date
Residence	
Cambridge, UK CB1 9YT	
Citizenship	
Great Britain	
Post Office Address	
3 Teasel Way	
Cambridge, UK CB1 9YT	